

# Infection prevention and control in the context of coronavirus disease (COVID-19): a living guideline

13 January 2023



World Health  
Organization

Preparedness  
IPC  
Response Readiness

**Contact**

WHO Health Emergencies - Infection Prevention and Control  
Avenue Appia 20, 1211 Geneva 27, Switzerland  
WHEIPC@who.int

**Infection prevention and control in the context of coronavirus disease (COVID-19): a living guideline, 13 January 2023**

© World Health Organization 2023. Some rights reserved. This work is available under the [CC BY-NC-SA 3.0 IGO](#) licence.

WHO reference number: WHO/2019-nCoV/ipc/guideline/2023.1

## Sections

Executive Summary .....	5
Definitions .....	6
Abbreviations .....	8
Methodology .....	9
Part 1: Health care settings .....	12
What is an IPC programme? .....	12
Environmental cleaning .....	13
Home care for patients .....	13
IPC when COVID-19 is suspected or confirmed .....	13
IPC principles and procedures for COVID-19 vaccination activities .....	13
Long term care facilities .....	13
Mask use .....	13
Mask use in health care setting acknowledgements, authorship, contributions .....	28
PPE Technical Specifications .....	29
Prevention, identification and management of health worker infection.....	30
Rational use of PPE and considerations during severe shortages .....	30
Risk assessment and management of exposure .....	30
Safe dead body management .....	30
Water, sanitation, hygiene, and waste management .....	30
Part 2: Community settings .....	31
Introduction to public health and social measures.....	31
Mask use .....	32
Mask use in the community .....	32
Type of mask used by the general public.....	41
Mask use during physical activity.....	45
Mask use in the community authorship, acknowledgements, and contributors .....	46
Mask use by children.....	48
Introduction .....	48
Age specific recommendations .....	51
Special populations .....	60
Implementation considerations for use of masks in schools .....	62
Mask use by children authorship, contributors and acknowledgements .....	62
Home care for patients.....	63
Water, sanitation, hygiene, and waste management .....	64
Safe dead body management .....	64

Annexes .....64

References.....68

## Executive Summary

Updated

### Version 4.0 Infection prevention and control in the context of coronavirus disease 2019 (COVID-19): a living guideline

#### Updated chapter: Mask use. Part 2: Community settings

#### About this guideline

The *Infection prevention and control in the context of coronavirus disease 2019 (COVID-19): a living guideline* consolidates infection prevention and control (IPC) technical guidance developed and published during the COVID-19 pandemic into evidence-informed recommendations for IPC. Part 1 presents IPC recommendations in the context of health care settings, while Part 2 presents recommendations for community settings. The methodology section describes the methodological approach used to develop the guideline, including a glossary. An annex is included with evidence tables for mask use in the health care and community settings and mask use by children. The *living guideline* is written, disseminated and updated on an online platform (MAGICapp). It has a user-friendly format and easy-to-navigate structure that accommodates dynamically updated evidence and recommendations. This structure focuses on what is new while keeping existing recommendations updated within the guideline.

This *living guideline* considers the current and evolving epidemiological trends for COVID-19 and the emergence of new variants of concern (VoC), including Omicron and its sublineages, and factors such as population immunity, availability and uptake of vaccines, and other contextual factors of the COVID-19 pandemic.

The target audiences of these guidelines are policy- and decision-makers, public health professionals, IPC professionals at the national, subnational and facility levels, health care facility administrators, managers, and other health workers.

#### Context

Countries are facing varying degrees of impact from severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). Considering the global COVID-19 epidemiological situation, the evolution and unpredictability of the virus, and the effect of current and future SARS-CoV-2 VoC, including descendent lineages of these variants and concurrent circulation of other respiratory viruses, including influenza, a combination of mitigation methods are required to reduce the impact of the COVID-19 pandemic [1]. Factors such as the intensity of SARS-CoV-2 circulation, vaccination coverage, population immunity to SARS-CoV-2, health systems' responsiveness capacity, and adjustability of public health and social measures (PHSM) influence viral circulation and epidemiological situations worldwide [3][4]. National policies should be strengthened and implemented to encourage the maintenance of IPC measures in healthcare facilities and PHSM for COVID-19 in community settings based on these and other factors [4][5]. In particular, prior to adjusting PHSM, the local situation should be determined, taking into consideration three main factors: SARS-CoV-2 transmissibility (also in consideration of any emerging variant); the seriousness of COVID-19; and the impact on the health system [65].

#### Updated recommendations

This version of the guidelines (version 4.0), includes the following recommendations:

- 1) a **strong recommendation** for mask use in community settings in higher risk situations;
- 2) a **conditional recommendation** to encourage a risk-based approach for mask use in community settings in situations that do not fall within the scope of the strong recommendation;
- 3) a **good practice statement** advising individuals with signs and symptoms suggestive of COVID-19 or those who test positive for COVID-19 to wear a medical mask when interacting with others in or outside of one's household or sharing space with others; and
- 4) amendment of an existing **good practice statement** on the implementation of policies to reduce SARS-CoV-2 transmission to promote the revisiting and strengthening of PHSM as new evidence arises.

#### Understanding the new recommendations

The Guideline Development Group (GDG) considered a combination of current scientific evidence [2] while assessing relative benefits and harms, values and preferences, resource implications, availability, and feasibility issues. The strong and conditional recommendations for mask use in the community complement each other, outlining possible scenarios in which masks may benefit the wearer. As some situations (see strong recommendation) necessitate mask wearing for maximum protection, others (see conditional recommendation) may be better suited for a risk-based approach. PHSM combine measures (for example, physical distancing, mask use, and hand hygiene) to mitigate the impact of SARS-CoV-2 infection.

### Updates to prior recommendations

Version 1.0 of the COVID-19 infection prevention and control living guideline: mask use in community settings published in December 2021 [6], provided new guidance on mask use in community settings. This guideline superseded existing advice in Mask use in the context of COVID-19 issued in December 2020 [7]. Version 4.0 of the *living guideline* is the most recent update on guidance for mask use in the community.

Version 2.0 of the COVID-19 living guideline jointly developed by the World Health Organization (WHO) and the United Nations Children's Fund (UNICEF) was published in March 2022 [8]. This version contained updated guidance on mask use in children; *Advice on the use of masks for children in the context of COVID-19* was first published in August 2020 as an annex to the document Mask use in the context of COVID-19 [7][9]. Version 2.0 of the *living guideline* superseded any previous advice on the use of masks for children in the context of COVID-19.

Version 3.0 of the COVID-19 living guideline published in April 2022 [10], provided new guidance on mask use in healthcare settings. This updated guideline superseded any advice on mask use in the health care setting in the context of COVID-19 [10][11][12].

### Guideline development

This guideline was developed using the Grading of Recommendations, Assessment, Development and Evaluation (GRADE) processes and Evidence to Decision framework [13] and in accordance with WHO norms and standards for guideline development [14][15].

The composition of the GDG includes experts in IPC, epidemiology, infectious diseases, paediatrics, water, sanitation and hygiene, engineering, aerobiology, and healthcare providers. The groups were balanced according to geographical and gender representation. Different GDGs were convened to address specific settings or populations (see authorship, contributors, and acknowledgements section). A methodologist with expertise in guideline development assisted the GDG in formulating the recommendations. While the GDG takes an individual patient perspective in making recommendations, it also considers resource implications, acceptability, feasibility, equity and human rights. The WHO Quality Assurance of Norms and Standards department helped to identify published rapid systematic reviews for the review process. Where required, WHO staff or commissioned external review teams conducted systematic reviews to address specific questions. Due to the rapidly evolving nature of the pandemic, preprints were included in the evidence synthesis. Additional details are described in the methodology section.

### Updates and access

This guideline and its previous versions are available through the WHO website and MAGICapp (online and PDF outputs for readers with limited internet access).

## Definitions

<i>Adequately ventilated patient room or area</i>	in health facilities where a mechanical ventilation system is available, the ventilation rate should be 6-12 air changes per hour (e.g., equivalent to 40-80 L/s/patient for a 4x2x3 m <sup>3</sup> room), and ideally 12 air changes per hour for new constructions, with a recommended negative pressure differential of $\geq 2.5\text{Pa}$ (0.01-inch water gauge) to ensure that air flows from the corridor into patient rooms" [16].
<i>Aerosol generating procedures (AGP)</i>	is a procedure that can induce the production of fine respiratory droplet in the patient[17]. AGPs are identified by the WHO as the following: tracheal intubation, non-invasive ventilation (e.g. bilevel positive airway pressure, continuous positive airway pressure), tracheotomy, cardiopulmonary resuscitation, manual ventilation before intubation, bronchoscopy, sputum induction by using nebulised hypertonic saline, dentistry and autopsy procedures. In oral health care, the following are considered AGPs: all clinical procedures that use spray generating equipment such as three-way air/water spray, dental cleaning with ultrasonic scaler and polishing; periodontal treatment with ultrasonic scaler; any kind of dental preparation with high or low speed hand pieces; direct and indirect restoration and polishing; definitive cementation of crown or bridge; mechanical endodontic treatment; surgical tooth extraction and implant placement. It remains unclear whether aerosols generated by nebuliser therapy or high-flow oxygen delivery are infectious or whether other procedures (e.g. nasogastric tube insertion, suctioning for airway clearance, or swabbing procedures) involve the risk of aerosol generation, due to lack of evidence or low quality evidence.
<i>Airborne transmission*</i>	is the spread of an infectious agent caused by the dissemination of droplet <i>nuclei</i> that remain infectious when suspended in air over long distances and time. Airborne transmission can be further categorized into obligate or preferential airborne transmission [18].

	<ul style="list-style-type: none"> <li>• Obligate airborne transmission refers to pathogens that are transmitted only by deposition of droplet <i>nuclei</i> under natural conditions (e.g. pulmonary tuberculosis) [18].</li> <li>• Preferential airborne transmission refers to pathogens that can initiate infection by multiple routes, but are predominantly transmitted by droplet <i>nuclei</i> (e.g. measles and chickenpox) [18].</li> <li>• Opportunistic airborne transmission refers to agents that naturally cause disease through other routes, but under special circumstances may be transmitted via fine particle aerosols [17].</li> </ul>
<i>A child</i>	any person under the age of 18 years [19].
<i>Contact transmission</i>	<p>is the spread of an infectious agent caused by physical contact of a susceptible host with people or objects.</p> <ul style="list-style-type: none"> <li>• Direct contact transmission involves both a direct body-surface-to-body-surface contact and physical transfer of microorganisms between an infected or colonized person and a susceptible host.</li> <li>• Indirect contact transmission involves contact of a susceptible host with a contaminated intermediate object (e.g. contaminated hands) that carries and transfers the microorganisms [18].</li> </ul>
<i>Droplet transmission</i>	<p>is the spread of an infectious agent caused by the dissemination of droplets. Droplets are primarily generated from an infected (source) person during coughing, sneezing and talking. Transmission occurs when these droplets that contain microorganisms are propelled (usually &lt; 1 m) through the air and deposited on the conjunctivae, mouth, nasal, throat or pharynx mucosa of another person. Most of the volume (&gt; 99%) comprises large droplets that travel short distances (&lt; 1 m) and do not remain suspended in the air. Thus, special air handling and ventilation are not required to prevent droplet transmission [18].</p>
<i>Hand hygiene</i>	is a general term that applies to handwashing, antiseptic handwashing, antiseptic hand rubbing or surgical hand antisepsis [18].
<i>Health care facility</i>	includes primary, secondary, tertiary care levels, outpatient care, and long-term care facilities.
<i>Health workers</i>	all people primarily engaged in actions with the primary intent of enhancing health. This includes health service providers, such as doctors, nursing and midwifery professionals, public health professionals, technicians (laboratory, health, medical, and non-medical), personal care workers, healers, and practitioners of traditional medicine. It also includes health management and support workers, such as cleaners, drivers, hospital administrators, district health managers, social workers and other occupational groups in health-related activities. This group includes those who work in acute care facilities and long-term care, public health, community-based care and other occupations in the health and social care sectors [20].
<i>Filtering facepiece respirators (FFR or respirators)</i>	<p>offer a balance of filtration, breathability and fit. Whereas medical masks filter 3-micrometre droplets, “N95” and “FFP2” rated FFRs must filter a more challenging 0.075-micrometre particles or particulates and do so across the entire surface of the respirator as a result of the fitted design. European “FFP2” FFRs, according to EN 149 standard, filter at least 94% Sodium Chloride (NaCl) salt particles and paraffin oil droplets. The United States of America “N95” FFRs, according to National Institute for Occupational Safety and Health (NIOSH) NIOSH 42 CFR Part 84, filter at least 95% NaCl salt particles. Certified FFRs must ensure unhindered breathing by meeting inhalation and exhalation breathing resistances below the maximum thresholds. Another important difference between FFRs and other masks is how filtration is tested. Medical mask filtration is assessed by testing filtration over a cross-section of the masks. In contrast, FFRs are tested for filtration across the entire surface. Most importantly, “FFP2” FFRs are fit-tested on a sample of human participants and the FFRs are measured for leaks as part of product certification. Similarly, for “N95” FFRs, individual workers are fit tested for specific FFRs at the workplace and typically on an annual basis. Therefore, in both cases, by ensuring the outer edges of the FFR seal around the wearer’s face, the FFRs filtration is closer to the actual filtration of inhaled air. Other FFR performance requirements include being within specified parameters for maximum CO2 build-up [21].</p>
<i>Medical masks</i>	<p>surgical or procedure masks that are flat or pleated and are affixed to the head with straps around the ears, the head or both. Their <u>performance standards</u> are tested according to a set of standardised test methods (American Society for Testing Materials (ASTM) ASTM F2100, EN 14683, or equivalent) that aim to balance high filtration, adequate breathability and, optionally, fluid penetration resistance [21].</p>

<i>Non-medical masks</i>	a type of facial covering of the mouth and nose of the wearer used to mitigate the spread of respiratory infections which does not meet the performance standards of 'medical' or 'surgical' masks. Their primary purpose is for source control and to provide a degree of particulate filtration to reduce the amount of inhaled particulate matter. Essential parameters for the performance and safety of non-medical masks have been advocated during the COVID-19 Public Health Emergency of International Concern (PHEIC) through several existing international guidelines and one international standard for non-medical masks (ASTM F3502-21) [21][22][23][24][25]. Non-medical masks which are self-made or commercially produced and do not meet guideline supported essential parameters are permitted in areas which have not mandated minimum performance requirements for non-medical masks prior to sale and for use by the general public.
<i>Standard precautions</i>	are routine IPC precautions that should apply to all patients, in all settings. They are intended to minimize the spread of infection associated with health care and to avoid direct contact with patients blood, body fluids, secretions and, non-intact skin. These precautions include hand hygiene, use of PPE based on risk assessment, respiratory hygiene, environmental cleaning and disinfection, waste management, reprocessing of medical devices, linen and laundry, management, prevention of needle-stick or sharps injuries [26][27].
<i>Universal masking</i>	the requirement for all persons (staff, patients, visitors, service providers and others) in health facilities to wear a mask at all times except when eating or drinking.
<i>Targeted continuous medical mask use</i>	the practice of wearing a medical mask by all health workers and caregivers working in clinical areas during all routine activities throughout the entire shift.
<i>Transmission based precautions</i>	are used in addition to standard precautions for patients who may be infected or colonized with certain infectious agents. These precautions include contact, droplet, and airborne precautions and should be implemented to prevent infection transmission [26][28].

\*Definition from the WHO Guidelines on "[Infection prevention and control of epidemic-and pandemic-prone acute respiratory infections in health care](#)" (2014) [18]. WHO has hosted expert global consultations in 2022 to further review and update the definition of airborne transmission: the results will be published in 2023. For the latest information on how COVID-19 is transmitted, please see "[Coronavirus disease \(COVID-19\): How is it transmitted?](#)".

## Abbreviations

AGP	Aerosol generating procedure
ASTM	American Society for Testing Materials
aOR	Adjusted odds ratio
COVID-19	Coronavirus disease 2019
CI	Confidence interval
CT	Community transmission
DOI	Declaration of interest
EtD	Evidence to decision
FFR	Filtering facepiece respirator
GDG	Guideline Development Group
GPS	Good practice statement
GRADE	Grading of Recommendations, Assessment, Development and Evaluation
HAI	Healthcare associated infection

ILI	Influenza-like illness
IPA	International Paediatric Association
IPC	Infection prevention and control
NIOSH	National Institute for Occupational Safety and Health
MAGIC	Magic Evidence Ecosystem Foundation
OR	Odds ratio
PICO	Population, intervention, comparator, outcome
PHEIC	Public Health Emergency of International Concern
PHSM	Public health and social measures
PPE	Personal protective equipment
RCT	Randomized control trial
SARS-CoV-1	Severe acute respiratory syndrome coronavirus
SARS-CoV-2	Severe acute respiratory syndrome coronavirus 2
UNICEF	United Nations Children's Fund
US CDC	United States Centers for Disease Control and Prevention
WHO	World Health Organization
VE	Vaccine effectiveness
VOC	Variant of concern

## Methodology

### Guideline Development Groups (GDG) and External Review Groups

The GDG was convened to review the available evidence and determine IPC recommendations, good practice statements (GPS) and implementation considerations included in this document. The GDG consisted of individuals with broad expertise spanning multiple specialties, across all WHO regions, and was gender balanced. Consensus was sought for recommendations and GPS. When consensus was not achieved, approval of a recommendation or GPS required a majority ( $\geq 70\%$ ) of the GDG voting members.

The responsible technical officer collected the required declaration of interest (DOI) from GDG members and assessed them for any potential conflicts. If a conflict of interest is identified, appropriate actions are taken in accordance with the [WHO Handbook for guideline development](#) and [WHO Guidelines for DOI \(for WHO Experts\)](#) [14][15]. This included removal from the GDG or recusal from voting or discussion for a particular recommendation or a decision to take no action.

External review group members were also identified for specific technical areas and provided additional review of the guidelines. External review groups do not change the recommendations made by the GDG; however, any major concerns are brought back to the GDG for additional discussion. For more information on authorship, contributions, and DOI, please refer to the acknowledgement section.

### Evidence synthesis and assessment

Given the dynamic situation of the COVID-19 pandemic, this living guideline integrates existing guidance that was developed using streamlined processes. As noted in the Executive Summary, with support from the WHO Quality Assurance of Norms and Standards department, rapid systematic reviews of published literature are identified for review. Due to the time lag for peer-reviewed publication of relevant studies in the context of a dynamic pandemic, preprints are included in the evidence synthesis. In addition, for some topics, systematic reviews are commissioned to external groups (clinical effectiveness of mask use in health care and community settings) or conducted by WHO staff (ecological studies on masks effectiveness). These reviews have been published and detail a search strategy within

the publication and are regularly updated to identify any emerging evidence that may inform deliberations by the GDG [2][29][30][31][32][33].

Evidence from randomized control trials (RCT) has been limited. Therefore, the reviews include non-randomized studies, cohort, case-control, and ecological studies. The systematic reviews are presented in GDG meetings and are supplemented by other (non-systematically reviewed) data presented by WHO staff, Member States, or partner organizations. Such presentations inform considerations regarding contextual factors on mask recommendations, mask filtration properties, and technical specifications on ventilation, including mask values/preferences, acceptability and feasibility, in the context of the changing epidemiology of COVID-19.

The literature for each identified topic is assessed using Grading of Recommendations, Assessment, Development and Evaluation (GRADE) to determine the certainty of the evidence (Table 1) based on the presence of risk of bias/study limitations, inconsistency, imprecision, indirectness and publication/reporting biases.

**Table 1. Determining the Quality of Evidence in Grading of Recommendations, Assessment, Development and Evaluation (GRADE)**

Quality level	Definition
<b>High</b>	The Group is very confident in the estimate of effect and considers that further research is very unlikely to change this confidence.
<b>Moderate</b>	The Group has moderate confidence in the estimate of effect and considers that further research is likely to have an important impact on that confidence and may change the estimate.
<b>Low</b>	The Group has low confidence in the estimate of effect and considers that further research is very likely to have an important impact on that confidence and is likely to change the estimate.
<b>Very low</b>	The Group is very uncertain about the estimate of the effect.

### Process for developing recommendations

Once the certainty of the evidence is determined, the GDG, with the guidance of the Methodologist, determines if a recommendation (strong or conditional) or a GPS is warranted. GRADE evidence profiles contain an assessment of the certainty of the evidence and a summary of findings for each critical outcome and each key question. The GDG uses these summaries as the basis for discussions and formulation of recommendations.

The Evidence to Decision (EtD) framework was used by the GDG to support the formulation of the recommendation or GPS. Core domains in the EtD framework are the balance of benefits and harms and the quality of the evidence, although other EtD domains (values/preferences, acceptability, feasibility, costs, and equity) also informed the recommendations (Table 2). The EtD framework for mask use in children was informed by five consultation sessions conducted by the United Nations Children's Fund (UNICEF) with members of the International Paediatric Association (IPA), and members from different geographical regions, in multiple languages, regarding paediatric health professionals' children's field experiences (including acceptability and feasibility) with the implementation of previous WHO guidance on masks. For other recommendations on mask use in health workers, the EtD domains were informed by invited presentations from stakeholders from individual countries regarding acceptability and feasibility, and by an invited presentation regarding mask availability and costs globally. The GDG also received regular updates on SARS-CoV-2 epidemiology and transmission from the WHO epidemiology team. Otherwise, the EtD domain assessments (including values/preferences and equity) were based on the collective input and experience of the GDG, which comprised members (including persons in the community, clinicians, and policymakers) who represented all WHO regions and ranged from low to very high income countries, supplemented by key studies suggested by GDG members when available. Additional systematic reviews were not commissioned and formal surveys outside of the GDG were not conducted to inform the EtD assessments further.

The GDG graded recommendations as strong or conditional. Strong recommendations are supported when benefits clearly outweigh harms with at least moderate certainty; other factors that support strong recommendations are non-sensitivity to variability in preferences/values regarding outcomes, wide feasibility and acceptability, cost savings or cost effectiveness, and likely positive impacts on improving equity. When certainty is low or very low, strong recommendations require a strong rationale for potential net benefits despite the limitations in the evidence and strong support from the other EtD domains. In these situations, GPS may be considered if the certainty of benefits is high based on indirect evidence (see the section on GPS). In some cases, after determining that the benefits of intervention do not outweigh the harms and considering EtD domains (Table 2), the GDG may make a recommendation against an intervention. The GRADE tables used in this living

guideline can be found in the Annex section of this living guidance.

The GRADE tables used in this living guideline can be found in the Annex section of this living guidance.

**Table 2. Evidence to Decision (EtD) framework**

Domain	Favours strong recommendations	Favours conditional recommendations
Balance of benefits and harms	Benefits highly outweigh harms	Benefits and harms more closely balanced
Quality of evidence	Higher certainty	Lower certainty
Values/preferences regarding outcomes	Benefits to harms assessment not impacted by variability in values/preferences	Variability in values/preferences would impact benefits to harms assessment
Acceptability	Highly acceptable	Low or variable acceptability
Costs/resources	Cost saving/cost effective	Costly/cost ineffective
Feasibility	Feasible in intended settings	Unfeasible or feasibility varies in intended settings
Equity	Increased equity	Decreased equity or effect on equity variable

**Good practice statements and implementation considerations**

GPS are most suitable when benefits are large and harm very small; the certainty of benefits and harms are great; the values and preferences are clear; the intervention is cost saving; and the intervention is clearly acceptable, feasible, and promotes equity. GPS characteristically represent situations in which a large and compelling body of indirect evidence, made up of linked evidence including several indirect comparisons, strongly supports the net benefit of the recommended action. GPS are generally issued due to various reasons, including the process, priorities, timeline, resources or nature of the evidence being assessed but are rooted in the fact that answers are obvious. GPS are not "GRADEd" statements [34].

Implementation considerations are critical elements that facilitate the appropriate use of recommendations and GPS but are not assessed using the GRADE methodology. They may be actionable and relevant to implementing one of the intervention options and may include information to enhance the implementation of the intervention [35].

**Readership cues for statements**

Table 3 presents the readership cues used for the statements in this living guideline. The green checkmark and red X symbols reflect statements that are developed using the GRADE evidence assessment methodology and the use of the evidence to decision framework to inform a recommendation or a GPS. The grey bar refers to implementation considerations that support statements through practical advice and are the product of expert consensus.

**Table 3. Readership cues used for statements in the living guideline**

	<p>The GREEN checkmark symbol denotes a recommendation or a good practice statement in favour of an intervention.</p>
---	---



The RED X denotes a recommendation or good practice statement against an intervention.

Implementation  
Consideration

The GREY bar denotes an implementation consideration supporting the practical implementation of the statement.

### Periodicity of the guideline revision and updates

Ongoing reviews are being conducted by WHO staff, as is the external living systematic review that has been commissioned to continuously monitor emerging evidence on the use of masks in the context of the COVID-19 pandemic. New evidence identified in these reviews that could inform revised or new recommendations will trigger reconsideration of the evidence by the GDG. Furthermore, as the pandemic evolves, including changes in transmission intensity, circulation of new variants of concern, and health systems' capacity to respond to new epidemiological scenarios, the GDG will review the current evidence on IPC and PHSM.

## Part 1: Health care settings

The document "[Infection prevention and control in the context of coronavirus disease \(COVID-19\): A living guideline](#)" brings together IPC technical guidance developed and published since the beginning of the COVID-19 pandemic. This consolidated document aims to provide users with the latest evidence-informed recommendations, through the MAGICapp platform, as a way to easily navigate guidelines in the dynamic context of COVID-19. Many parts of the technical guidance related to *Part 1: Health care settings* are currently under review. Links to the most recent publication of the technical guidance are available in the sections that follow. Updated guidelines on health care settings will be available in this living guideline in the near future.

## What is an IPC programme?

### What is an Infection Prevention and Control Programme?

Infection prevention and control (IPC) is a practical, evidence informed approach to preventing patients and health workers from being harmed by avoidable infections. Healthcare-associated infections (HAI) are among the most common adverse events in care delivery and a major public health problem impacting morbidity, mortality and quality of life. On average, 7% of patients in developed and 15% in developing countries will acquire at least one HAI at any one time [36]. These infections also present a significant economic burden at the societal level. However, a large percentage are preventable through effective IPC measures.

### Establishing an infection prevention and control programme at national and acute health care facility levels

The *WHO Guidelines on core components of infection prevention and control programmes at national and acute health care facility levels* [37] are the foundation of WHO strategies to prevent current and future threats from infection and antimicrobial resistance in health care. The core components constitute a framework of recommendations of good practices statement distributed into eight areas: 1) infection prevention and control programmes, 2) national and facility-level infection prevention and control guidelines, 3) infection prevention and control education and training, 4) healthcare-associated infections surveillance, 5) multimodal strategies for implementing infection prevention and control activities, 6) monitoring and evaluation and feedback, 7) workload, staffing and bed occupancy at the facility level and, 8) built environment, materials and equipment for infection prevention and control at the facility level. Ensuring adequate clinical staffing levels is recommended as a core component to prevent the transmission of HAI and multidrug-resistant organisms (MDRO), limit human-to-human transmission, reduce secondary infections, and prevent the transmission through amplification and super spreading events.

Considering that implementing an Infection Prevention and Control Programme requires a stepwise approach [38][39] to its full achievement, minimum requirements [40] have been identified to support it in countries where IPC is limited or nonexistent. In this regard, a facility level IPC programme with a dedicated and trained IPC team, or at minimum, an IPC focal point, should be in place and supported by the national and facility senior management. Achieving the IPC minimum requirements (and more robust and comprehensive IPC programmes in all countries is essential to sustain efforts to control the COVID-19 pandemic, other emerging and re-emerging pathogens, and multi drug resistant organisms (MDRO). Finally, WHO has also developed guidance on the core

competencies [41] required for infection prevention and control professional staff, which can be used for developing curricula for IPC specialists.

## Environmental cleaning

The most up-to-date technical guidance for "[Cleaning and disinfection of environmental surfaces in the context of COVID-19: interim guidance](#)" was published 15 May 2020. This guidance is under review and is pending integration into "[Infection prevention and control in the context of coronavirus disease \(COVID-19\): A living guideline](#)".

## Home care for patients

The most up-to-date guidance for "[Home care for patients with suspected or confirmed COVID-19 and management of their contacts: interim guidance](#)" was published 12 August 2020. This guidance is under review and is pending integration into "[Infection prevention and control in the context of coronavirus disease \(COVID-19\): A living guideline](#)".

## IPC when COVID-19 is suspected or confirmed

The most up-to-date technical guidance for "[Infection prevention and control during health care when coronavirus disease \(COVID-19\) is suspected or confirmed: interim guidance](#)" was published 12 July 2021. This guidance is under review and is pending integration into "[Infection prevention and control in the context of coronavirus disease \(COVID-19\): A living guideline](#)".

## IPC principles and procedures for COVID-19 vaccination activities

The most up-to-date technical guidance for "Aide-memoire: infection prevention and control (IPC) principles and procedures for COVID-19 vaccination activities" was published 15 January 2021. This guidance is under review and is pending integration into "[Infection prevention and control in the context of coronavirus disease \(COVID-19\): A living guideline](#)".

## Long term care facilities

The most up-to-date guidance for "[Infection prevention and control guidance for long-term care facilities in the context of COVID-19: interim guidance](#)" was published 21 March 2020. This guidance is under review and is pending integration into "[Infection prevention and control in the context of coronavirus disease \(COVID-19\): A living guideline](#)".

## Mask use

## Background

The WHO continuously reviews available data on SARS-CoV-2 variants of concern. For this version, the global epidemiological situation of the COVID-19 pandemic as of 21 January 2022 – at a time when the Omicron VOC had been identified in 171 countries across all six WHO Regions and was rapidly replacing Delta worldwide – was considered [42].

Omicron has a substantial growth advantage, higher secondary attack rates and a higher observed reproduction number than Delta. There is now significant evidence that immune evasion contributes to the rapid spread of Omicron. Other factors may be a shorter serial interval (by about 0.8 to 1.2 days compared to Delta) and potential increased intrinsic transmission fitness [42]. There is growing evidence that with Omicron, there is lower vaccine effectiveness (VE) against infection and symptomatic disease soon after vaccination compared to Delta. There is also evidence of accelerated waning of VE over time of the primary series against infection and symptomatic disease for the studied vaccines. Further studies are required to better understand the drivers of transmission and declining incidence in various settings. These factors include the intrinsic transmission fitness properties of the virus, degree of immune evasion, vaccination coverage and level of vaccine-derived and post-infection immunity, levels of social mixing and degree of application of public health and social measures (PHSM).

Essential measures to prevent SARS-CoV-2 transmission in health care facilities remain valid in the context of Omicron and should be strengthened [42].

WHO recommends using face protection as part of a comprehensive package of prevention and control measures to limit the spread of SARS-CoV-2. National policies and health facilities must continue to achieve and maintain IPC measures, including having an IPC programme or at minimum a dedicated and trained IPC focal point in place. Other necessary measures include engineering, environmental and administrative controls, standard and transmission based-precautions, screening and triage for early identification of cases and COVID-19 surveillance and vaccination of health workers. This is particularly important considering the rapid spread of Omicron and the high proportion of individuals who may be infected but are asymptomatic [42].

This document guides decision makers and IPC professionals to develop and implement policies on mask use in health care settings.

---

*Published 25 April 2022.*

## In areas of known or suspected community or cluster transmission

Strong recommendation for , Very low certainty evidence



### **In areas of known or suspected community or cluster SARS-CoV-2 transmission, universal masking is recommended in health care facilities:**

- In settings where caring for non-COVID-19 patients, unless differently specified (e.g. AGP), all health workers, including community health workers and caregivers, other staff, visitors, outpatients and service providers, should wear a well-fitting medical mask at all times within the health facility and in any common area (e.g., cafeteria, staff rooms).
- Inpatients are not required to wear a medical mask unless physical distancing of at least 1 metre cannot be maintained (e.g., during examinations or bedside visits) or when outside of their care area (e.g., when being transported), provided the patient is able to tolerate the mask and there are no contraindications.
- [Click here](#) for the recommendation on the mask type for health workers when caring for a suspected or confirmed COVID-19 patient.

---

*Published 25 April 2022.*

## Practical Info

When adopting universal masking within a health facility, it is essential health workers follow proper mask-wearing procedures and practices. For additional information review the implementation considerations on [mask management for health workers](#).

The [WHO recommendation on mask fitting](#) should be followed, including the related considerations on this critical aspect.

## Evidence To Decision

### Benefits and harms

Substantial net benefits of the recommended alternative

The wearing of a medical mask is associated with a decreased risk of acquiring SARS-CoV-2 infection [30]. In areas where there is community transmission of COVID-19, universal masking has been adopted by most hospitals to reduce potential transmission between health workers and other staff, patients, and those entering the facility. Five studies found that implementing a universal masking policy in hospital systems was associated with decreased risk of healthcare-acquired SARS-CoV-2 infection [43][44][45][46][47]. However, these studies have limitations, as most originated in the United States of America, and used a before-after design. Other limitations include lack of or limited control for confounders, such as the use of other personal protective equipment and exposures [43][44][45][46][47]. Furthermore, potential sensitivity to time periods selected for analysis for pre-and post-universal masking was identified, but none of the studies included sensitivity analysis. Literature provides limited insight into the harms of universal masking; evidence on mask use, in general, indicates bothersome but non-serious harms. Therefore, despite the limitations in the evidence, the GDG judged that the benefits of implementing universal mask use in healthcare facilities outweigh potential harms.

### Certainty of the Evidence

Very low

Given the limited number and the type of evidence available (i.e., before-after studies) regarding the implementation of universal masking as an IPC procedure, the certainty of the evidence is rated as very low. However, despite the very low certainty of evidence pertaining to universal masking, the wearing of a medical mask is associated with a decreased risk of acquiring SARS-CoV-2 infection [30].

### Values and preferences

No substantial variability expected

Given the protective effects associated with mask use, health workers, including community health workers and caregivers, would likely favour the implementation of universal masking [30][43][44][45][46][47][47]. In the context of universal masking, some health workers may prefer to wear respirators instead of a medical mask, based on their perception of what offers the better protection to prevent SARS-CoV-2 infection and emergent evidence that the use of respirators might be more effective in the control of transmission of some variants of concern such as Omicron. There are no substantial variabilities in values and preferences.

### Resources

Implementing universal masking is likely to have a low to moderate impact on resources.

### Equity

No important issues with the recommended alternative

No adverse impacts on equity to the individual have been identified, as long as masks are provided by health care facilities and are readily available for all health workers, staff, visitors and patients.

### Acceptability

No important issues with the recommended alternative

Universal mask use is likely to be easily accepted in health care facilities given the protective effects for health workers, other staff, visitors and patients [30][43][44][45][46][47][47].

## Feasibility

No important issues with the recommended alternative

The universal use of masks in health care facilities is likely feasible and is currently the standard in most countries, in the context of the COVID-19 pandemic.

## Justification

Upon deliberations during the GDG meeting, the decision regarding this recommendation's strength was reached through online voting. Despite the very low certainty of the evidence for the implementation of universal masking, the evidence does indicate benefits without significant harms; in addition, the GDG members judged that universal masking could prevent potential serious harms of health care worker infections and transmission in health care. Of 28 members of the GDG, 78.6% (22) voted that this should be a strong recommendation. Members also felt that based on their own professional experience, or that of colleagues, universal masking in health settings is already routine in most countries; therefore, the acceptability and feasibility favoured a strong recommendation, as well. Furthermore, the utilization of a medical mask is associated with a decrease in SARS-CoV-2 transmission<sup>[31]</sup>.

Additionally, the GDG reviewed the mask type to be used universally in health care facilities. In light of new VOCs with increased transmissibility and the subsequent need to better protect health workers and their patients, GDG members felt the exclusive use of medical masks was justified. Given the available evidence on mask effectiveness of medical masks and their requirement to adhere to strict standards, a majority of members felt the universal use of medical masks in the health care setting would provide better protection for staff, patients, visitors and the community.

## In areas of known or suspected sporadic transmission

Conditional recommendation for , Very low certainty evidence



**In areas of known or suspected sporadic SARS-CoV-2 transmission, targeted continuous medical mask use is recommended in health care facilities:**

- In settings when caring for non-COVID-19 patients, health workers, including community health workers and caregivers who work in clinical areas, should continuously wear a well-fitting medical mask during routine activities throughout the entire shift, unless differently specified (e.g. when performing AGP) and apart from when eating and drinking.
- In non-patient areas, staff are not required to wear a medical mask during routine activities if they have no patient contact.
- [Click here](#) for the recommendation on mask type for health workers when caring for a suspected or confirmed COVID-19 patient.

*Published 25 April 2022.*

## Practical Info

When adopting targeted continuous masking within a health facility, it is essential health workers follow proper mask-wearing procedures and practices. For additional information review the implementation considerations on [mask management for health workers](#).

The [WHO recommendation on mask fitting](#) should be followed, including the related considerations on this critical aspect.

## Evidence To Decision

### Benefits and harms

Substantial net benefits of the recommended alternative

Given the protective effects of mask use, the benefits of implementing targeted continuous mask use in healthcare facilities outweigh potential harms <sup>[30]</sup>. Five studies found that consistent mask use in health care facilities was associated with a

decreased risk of SARS-CoV-2 infections in health workers. However, it is essential to note that these studies only investigated universal masking, not targeted continuous masking. The effects found in these studies have been extrapolated for the aforementioned recommendation [43][44][45][46][47].

### Certainty of the Evidence

Very low

The evidence for targeted continuous masking has been extrapolated from evidence on universal masking; therefore, the certainty of the evidence is rated as very low. However, despite the very low certainty of evidence pertaining to targeted continuous masking, the wearing of a medical mask is associated with a decreased risk of acquiring SARS-CoV-2 infection [30].

### Values and preferences

No substantial variability expected

Given the protective effects of mask use, health workers, including community health workers and caregivers, would likely favour targeted continuous masking in health facilities [30][43][44][45][46][47]. There are no important variations in the values and preferences.

### Resources

No important issues with the recommended alternative

The implementation of targeted continuous masking is likely to have a low to moderate impact on resources.

### Equity

No important issues with the recommended alternative

This intervention will likely cause no adverse impacts on equity, so long as masks are provided in health care settings and are readily available.

### Acceptability

No important issues with the recommended alternative

The universal use of masks in healthcare facilities is the standard in most countries in the context of the COVID-19 pandemic and has been widely implemented.

### Feasibility

No important issues with the recommended alternative

The use of targeted continuous mask use in health care facilities is likely feasible.

## Justification

After GDG members discussed their perspectives on recommending the implementation of targeted continuous masking, the decision to formalise the above statement as a conditional recommendation was reached through online voting. GDG members felt a conditional recommendation was well suited for this guidance, given that the evidence for continuous masking was inferred from evidence on universal masking and the statement is written for an epidemiological situation with few COVID-19 cases.

### Implementation consideration

The following procedures and practices should be ensured when wearing a mask in health care settings [48]

- Medical masks should be combined with other measures including frequent hand hygiene and physical distancing of at least 1 metre among health workers in shared and crowded places such as cafeterias, break rooms and dressing rooms [49].
- Medical masks must be changed when wet, soiled or damaged or if the health worker or caregiver removes the mask for any reason (e.g. for eating or drinking or caring for a patient who requires droplet/contact precautions for reasons other than COVID-19).
- Used medical masks should be disposed of properly.
- The medical mask should not be touched to adjust it or if it is displaced from the face for any reason. If this happens, the mask should be safely removed and replaced and hand hygiene performed.
- The medical mask (as well as other PPE) should be discarded and changed after caring for any patient who requires contact/droplet precautions for other pathogens, followed by hand hygiene.
- Under no circumstances should a medical mask be shared between health workers.
- Medical masks can become displaced from their optimal placement, over the mouth and nose, during extended use, which creates gaps for respiratory particles to bypass the filtration layers on inhalation and exhalation [50]. The [WHO recommendation on mask fitting](#) should be followed, including the related considerations on this critical aspect.

Published 25 April 2022.

### In any transmission scenario

#### Good practice statement



Appropriate mask fitting should always be ensured (for respirators, through fit testing and a user seal check when a filtering facepiece respirator is donned; and for medical masks, through methods to reduce air leakage around the mask) as well as compliance with appropriate use of PPE and other standard and transmission-based precautions.

Published 25 April 2022.

### Practical Info

#### Implementation consideration

#### Methods to improve the fit of respirators or medical masks

##### **Respirators**

- Filtering facepiece respirators (FFRs) vary for their measurement of fit, either through maximum allowable leak tightness or minimum fit factor. For European certified FFRs, the maximum leakage varies from:
  - FFP1 (maximum 22% leakage)
  - FFP2 (maximum 8% leakage) and
  - FFP3 (maximum 2% leakage)
  - European certified FFRs (EN 149) are subject to testing for leakage with human participants as part of the product's certification.

- For NIOSH, N-type FFRs (minimum fit factor of 100) are certified according to OSHA 29 CFR 1910.134 for each wearer prior to use.
- At a minimum, FFRs that meet FFP2 and N95 performance levels are recommended to be worn by health workers in areas where AGP are performed [21].
- Ensure a range of FFR sizes are available to accommodate different face shapes and sizes, especially for those with small faces.
- Qualitative or quantitative fit testing should be performed annually and for new staff at the employer's expense to ensure that the respirator model fits each health worker's unique facial features and provides a consistent seal [51].
- A seal check should be performed on FFRs whenever donned by a health worker to determine if the adequate fit is achieved by the specific FFR they have donned. See [WHO guidance on how to perform a particulate respirator seal check](#) for additional details.

**Two methods can be used for fit testing FFRs**

1) qualitative fit test (health worker reports taste of an ambient aerosol) and 2) quantitative fit test

	Qualitative Fit Testing	Quantitative Fit Testing
<b>Standard test methods</b>	OSHA 29 CFR 1910.134 Appendix A (for N95)	EN 149, Clause 7.9.1 (EN-type, e.g., FFP2) OSHA 29 CFR 1910.134 Appendix A (e.g., N95)
<b>Equipment</b>	Hood and sweet/bitter aerosol	Ambient aerosol condensation nuclei counter
<b>Pass/Fail Criteria</b>	Wearer report tasting aerosol	>8% leakage (for FFP2) <100 fit factor (for N95)

**Medical masks**

Improving the fit of medical masks may not always be possible in low resource settings, given the resource requirements. However, techniques such as the “tie-and-tuck” method may benefit low- and middle-income countries since they do not require additional materials. The “knot-and-tuck” and “linking-ear-loops-behind-the-head” techniques improve medical mask fit by reducing gaps on the sides of medical masks with ear loops. Such gaps allow air leakage (potentially containing infectious particles) to bypass the filtration layers of the medical mask when the wearer inhales or exhales.

**Considerations on the use of linking-ear-loops-behind-the-head techniques to improve medical mask fit**

- Always use a clean, unused rectangular pleated medical mask meeting the minimum performance standards (or equivalent)[21].
- Always clean hands thoroughly (per [WHO guidance](#)) prior to donning, doffing and/or manipulating a mask.
- Where connectors are used to link ear loops behind the head, ensure that these connectors are clean for use upon donning (either new, cleaned and disinfected or laundered, depending on the connector and local implementation strategy). When connectors are doffed, they should be treated as potentially contaminated. A local strategy should be in place to manage used connectors through cleaning and disinfection processes, laundering or discarding used connectors through standard waste management.

# How to improve medical mask fit in health care settings




**When linking ear loops behind the head**



**When using knot-and-tuck method**



Clean hands thoroughly before putting on and before and after taking off your mask



Attach a clean connector to link ear loops together\*



Fold the mask horizontally



Place the medical mask colour-side facing outward, attach ear loops behind ears



Make a knot on both ear loops as close to the edge of the mask as possible



Attach ear loops using connector behind head tightly



Push the extra material under the mask inward to ensure no gaps on both sides



Adjust the wire at the bridge of the nose and ensure there are no gaps between the mask and your face at the sides of your nose, cheeks, and under your chin.

\*Find a clean practical connector to link your ear loops, it can be:



adjustable rope



silicone

\*\*If a surface is used to fold and manipulate the mask, clean the surface first using a cloth wipe with soap and water, followed by disinfection using a cloth wipe soaked in 70-90% alcohol OR 0.1% sodium hypochlorite (or comparable hospital grade disinfectant) and allow for least 1 minute contact time before surface is used.

When adopting a mask policy, it is essential health workers follow proper mask-wearing procedures and practices. For additional information review the implementation considerations on [mask management for health workers](#).

### Justification

GDG members were asked if WHO should consider developing practical advice on improving medical mask fit, where's the majority of GDG members agreed this would be useful. Five options to improve the fit of masks were presented: the "use of ear loops linked behind the head"; the "tie-and-tuck method"; the use of a brace/fitter; the use of masks with ties behind the head instead of ear loops; double masking.

Twenty-five GDG members (83.3%) agreed with the use of ear loops linked behind the head; 23 (76.7%) agreed with the use of the tie-and-tuck method; 19 (63.3%) agreed with the use of masks with ties behind the head instead of ear loops; 16 (53.3%) agreed with the use of a brace/fitter; 10 (33.3%) agreed with double masking. Therefore, the use of ear loops linked behind the head and of the tie-and-tuck method was retained as advisable methods to improve the fit of masks and additional details can be found in the practical information section.

GDG members reported that the evidence available on improving the fit of medical masks to reduce the transmission risk of SARS-CoV-2 is in the form of laboratory-based studies with limited field and clinical investigations.

Strong recommendation for , Very low certainty evidence



A respirator should always be worn along with other PPE\* by health workers performing aerosol-generating procedures (AGP) and by health workers on duty in settings where AGP are regularly performed on patients with suspected or confirmed COVID-19, such as intensive care units, semi-intensive care units or emergency departments.

\*PPE includes gown, gloves, eye protection.

Published 25 April 2022.

### Practical Info

When adopting a mask policy within a health facility, it is essential health workers follow proper mask-wearing procedures and practices. For additional information review the section on [mask management for health workers](#).

The [WHO recommendation on mask fitting](#) should be followed, including the related considerations on this critical aspect.

### Evidence To Decision

#### Benefits and harms

Small net benefit, or little difference between alternatives

Among health care workers, exposure to an AGP such as tracheal intubation was associated with a higher risk of infection with SARS-CoV-1, the most closely related human coronavirus to SARS-CoV-2 [52]. Furthermore, a living rapid review showed that certain exposures such as involvement in intubations are significantly associated with SARS-CoV-2 infections [32][33]. However, no specific evidence assessing the effectiveness of different types of masks to prevent transmission of SARS-CoV-2 during AGP is available. Indirect evidence from laboratory simulation data provides insight on the plausibility and viability of aerosolised SARS-CoV-2 [53].

Respirators have higher filtration efficiency standards and demonstrate better fit with fewer air gaps allowing bypass of the filter media than the most commonly used rectangular medical masks, provided that they are appropriately fit tested and worn. Therefore, respirators are likely to be superior in preventing transmission of SARS-CoV-2 during AGP [32][33].

### Certainty of the Evidence

Very low

Given the absence of direct evidence related to SARS-CoV-2 and the limitations of the indirect evidence, the certainty of the evidence for the utilization of particulate respirators for patients with suspected or confirmed COVID-19 during AGP was rated as very low.

### Values and preferences

No substantial variability expected

Health care workers would highly prefer to wear a respirator during AGP in order to benefit from the perception of a higher protective effect. Thus, no variability is expected in health care workers' preferences related to the use of respirators during AGP to prevent transmission risk in preventing transmission.

### Resources

Important issues, or potential issues not investigated

#### Resource implications

The use of respirators requires an additional investment of financial and logistical resources, including the need for fit testing for all staff, requiring additional investments and expertise [51]. Some clinical and operational challenges may be experienced, in particular in low and middle income countries, and investments are needed in order to provide the best protection possible during AGP.

#### Knowledge gaps, research needs and comments

Additional research is needed to clarify which medical procedures produce aerosols and thus, potentially increase the transmission risk of SARS-CoV-2 and other respiratory pathogens thus, leading to the need for a higher level of respiratory protection. Conducting trials to compare the effectiveness of different types of masks to prevent infection during AGP would be unethical.

### Equity

Important issues, or potential issues not investigated

Given the limited global supply of respirators and their high cost in particular for resource-limited settings, inequity issues likely exist.

### Acceptability

No important issues with the recommended alternative

Stakeholders and policymakers' will likely accept the recommended use of respirators during procedures that produce aerosols as this is the policy currently in place in most countries and historically integrated into a conditional recommendation by the WHO for acute respiratory infections [18].

### Feasibility

No important issues with the recommended alternative

The use of respirators during the performance of an AGP is feasible although some resources implications have been noted.

### Justification

A majority of GDG members noted that despite the very low certainty of evidence, the acceptability and feasibility of implementation and the benefits of wearing a respirator during the performance of an AGP on a suspected or confirmed COVID-19 patient justified a strong recommendation. The GDG agreed to upgrade the strength of this recommendation from a conditional recommendation to a strong recommendation [18]. The decision was made in light of the increased widespread transmission of Omicron, its immune escape, and still limited vaccination coverage in health care workers worldwide.

Conditional recommendation for , Very low certainty evidence



A respirator or a medical mask should be worn by health workers along with other PPE– a gown, gloves and eye protection – before entering a room where there is a patient with suspected or confirmed COVID-19.

Respirators should be worn in the following situations:

- In care settings where ventilation is known to be poor\* or cannot be assessed, or the ventilation system is not properly maintained
- Based on health workers' values and preferences and on their perception of what offers the highest protection possible to prevent SARS-CoV-2 infection.

\*Ventilation in a health care setting is considered to be poor when the requirements established for these settings are not in place (see "Definitions" section).

*Note: This recommendation applies to any setting where regular care is provided to patients with suspected or confirmed COVID-19, including home care, long-term care facilities and community care settings. For settings where AGP are regularly performed on patients with suspected or confirmed COVID-19, see the strong recommendation above.*

Published 25 April 2022.

## Practical Info

When adopting a mask policy within a health facility, it is essential health workers follow proper mask-wearing procedures and practices. For additional information review the implementation considerations on [mask management for health workers](#).

The [WHO recommendation on mask fitting](#) should be followed, including the related considerations on this critical aspect, like the type of FFR that should be used by health workers.

## Evidence To Decision

### Benefits and harms

Small net benefit, or little difference between alternatives

The recommendation noted above made no distinction between the use of medical masks and respirators when caring for a COVID-19 positive patient, except in situations where respirators are clearly needed (e.g., AGP). Respirators have higher filtration efficiency standards and demonstrate better fit with fewer air gaps allowing bypass of the filter media than the most commonly used rectangular medical masks, provided that the respirators are appropriately fit tested and properly worn [21][51]. Given the protective effects of respirators, several GDG members advised that respirators may be superior to medical masks in preventing SARS-CoV-2 infection and their use should be encouraged when the health care worker delivers care in close contact with the patient and/or when ventilation is inadequate.

Evidence comparing the effectiveness of respirators versus medical masks for SARS-CoV-2 in healthcare settings is limited to five observational studies [54][55][56][57][58], which were conducted prior to the emergence of the Delta, Omicron and other variants and before widespread vaccination in healthcare settings. These five observational studies had methodological limitations (for example, recall bias, low participation, limited measurement of exposures) and reported inconsistent findings regarding the risk of SARS-CoV-2 infection between the use of respirators versus medical masks. One study showed a reduction of risk with respirator use [55], while in another two studies the use of respirators was not significantly associated with risk reduction [57][58]. One study showed no association [58], and another found respirators were associated with increased risk (OR 7.1), likely related to confounding factors [56]. Prior randomised controlled trials comparing respirators versus medical masks for prevention of clinical influenza-like illness (ILI) found no difference [59][60][61][62][63]. Overall, the strength of this evidence was rated as insufficient to recommend one type of mask versus the other.

The following side effects have been reported with respirators: discomfort, headaches, possible development of facial skin lesions, irritant dermatitis or worsening acne when used frequently for long hours [30]. Medical masks are typically associated with less discomfort or side effects than respirators given decreased thickness and reduced seal, although this has not been quantified. Undesirable outcomes from the prolonged use of respirators were noted, including general discomfort, headaches and the development of facial skin lesions, irritant dermatitis or worsening acne [30]. The fitting process for respirators is burdensome, and issues with achieving it have been well described. Furthermore, other factors may influence the overall risk of transmission, including general PPE use, PPE training, fit testing, ventilation, and behavioural factors (including compliance) as well as the fact that transmission of SARS-CoV-2 among health workers appears to mostly occur in community settings [30]. The balance of desirable and undesirable outcome effects was rated as uncertain. It was deemed uncertain whether respirators are more effective than medical masks in settings without exposure to AGP.

### Certainty of the Evidence

Very low

Given the methodological limitations of the evidence, notably inconsistency and indirectness (e.g. most studies conducted before the emergence of the Delta variant and none in the Omicron era), evaluation of non-SARS-CoV-2 infections or assessment of non-clinical outcomes, [30] the certainty of the evidence for particulate respirators versus medical masks was rated as very low.

### Values and preferences

Substantial variability is expected or uncertain

There is substantial variability in preferences related to the use of respirators in preventing HAI. In the context of the increased transmissibility of the Delta or Omicron variant, some health care workers may value the wider use of respirators to potentially reduce their risk, despite the limited evidence, as a precautionary approach. Others may not prefer to wear a respirator throughout their shifts because of discomfort and potential side effects. Local values, preferences and practicalities should play an important role in directing choices on the use of respirators versus medical masks.

### Resources

Important issues, or potential issues not investigated

#### Resource implications

The use of respirators for the care of all patients with suspected or confirmed COVID-19 in health care facilities requires an additional investment of financial and logistical resources, which could be challenging, in particular, in low and middle income countries. There is also the need for fit testing for all staff, requiring additional investments and expertise; however, scaling up the market for respirators could lead to cost reduction.

#### Knowledge gaps, research needs and comments

Randomised controlled trials on respirators versus medical masks in health care settings are in progress. Well-conducted observational studies on respirators versus medical masks and the risk of SARS-CoV-2 infection in healthcare settings in the context of the Omicron and other variants are urgently needed. More research is also needed to investigate the risks associated with medical masks and respirators and adverse events (including self contamination) during extended and repeated use. Other gaps include studies on simpler, faster and less costly methods, or alternative methods, to determine respirator fit and seal. Further data is needed regarding compliance with appropriate PPE use, including masks, and in particular, appropriate donning and doffing practices in COVID-19 and non COVID-19 units.

### Equity

Important issues, or potential issues not investigated

Given the limited global supply of respirators and their higher cost compared to medical masks, a recommendation to use respirators for all COVID-19 cases in health care settings could result in inequity in resource limited settings. However, it is also

expected that the widespread use of respirators (if available) will reduce inequities related to COVID-19 exposure risk. Unvaccinated health care workers worldwide are still at higher risk for infection, sometimes resulting in severe disease and death. There is an additional equity issue around medical masks, which may also not be available in sufficient quantities and of adequate quality in low resource settings.

### Acceptability

No important issues with the recommended alternative

The current recommendation provides the option of using either respirators or medical masks, except for specific circumstances when a respirator is required. Given this flexibility, it should be acceptable for stakeholders' and policymakers'.

### Feasibility

Important issues, or potential issues not investigated

Although WHO unpublished modelling data indicated an inadequate supply of respirators to replace medical masks in all COVID-19 health care settings, policies advising respirators in all COVID-19 settings would likely lead to increased investments and production. Furthermore, a strong supply distribution and logistics system is needed to ensure efficient procurement and reach across the whole health system. However, inefficiencies in the distribution of supplies and supply chain problems have been reported. The adequate fit of the device is correlated with the effectiveness of the FFP, but fit testing may not be feasible in all regions.

## Justification

The GDG considered the very low certainty of the evidence for particulate respirators versus medical masks, and agreed that the strength of this evidence was insufficient to recommend one type of mask versus the other, except in some specific conditions. However, many GDG members saw relevance in the epidemiological evidence showing that the Omicron variant is spreading significantly faster than the Delta variant in countries with documented community transmission [42]. Serious concerns were expressed about the evidence of SARS-CoV-2 re-infection with Omicron, and the data showing a reduction in neutralising antibody titres against Omicron and a significant reduction in VE against infection and symptomatic disease for Omicron compared to Delta [42]. Some GDG members also highlighted the fact that as of November 2021, 65% of health workers in 135 countries were fully vaccinated, but the vaccination status was unknown for 77.7 million health workers (58% of the global health workforce) [64].

Following in-depth discussions, the GDG was asked to decide whether to maintain the recommendation on the type of mask to be used in COVID-19 settings included in the Annex to "Infection prevention and control during health care when coronavirus disease (COVID-19) is suspected or confirmed". Out of 33 IPC GDG members who voted, 24 (68%) would have preferred to maintain the previous recommendation on the type of mask to be used in COVID-19 settings. The previous recommendation took into strong consideration serious concerns about the limited availability of respirators in low and middle income countries and the resource implications of more widespread use of respirators. The GDG voting on this recommendation in light of Omicron was based on the very low certainty of the evidence for particulate respirators versus medical masks, given the methodological limitations of the evidence, as well as the previous concerns about respirators' availability. However, among GDG members who advised maintaining the previous recommendation, 48% also stated that they would consider it acceptable to recommend either respirators or medical masks; 33% of them even considered it acceptable to prioritise respirators; the remaining 19% of them stated that no other option but maintaining the previous recommendation would be acceptable to them. 32% of the GDG members voted against maintaining the previous recommendation; among these, 70% advised recommending either respirators or medical masks, whereas 30% recommended the sole use of respirators in all settings where COVID-19 patients are given care. Given the limitations described, the deliberations of the GDG and decision-making process were also informed by the perspectives and experience of experts represented in the panel.

Following very careful interpretation of the GDG considerations and voting results which showed that 68% of GDG members would have preferred to maintain the recommendation included in the [Annex](#) issued on 1 October 2021, but also indicated what changes would have been acceptable to these GDG members, WHO decided to make this new conditional recommendation which was issued as [rapid guidelines on 22 December 2021](#). Given the increased transmissibility and rapid spread of the Omicron VOC, WHO leadership felt it was necessary to take a precautionary approach, according to the hierarchy of controls, and add the option of respirators to the recommendation on masks to use when entering a room with a COVID-19 positive patient regardless of AGP

being performed, despite the limitations of the available evidence on respirators versus medical masks in health care facilities.

**Table 2. Mask use in health care settings depending on transmission scenario, target population, setting, activity and type\***

Transmission scenario	Target population	Setting	Activity	Mask type
Any transmission scenario	Health workers	Health care facility**	Performing an AGP or providing care in a setting where AGP are in place for suspected/confirmed COVID-19 patient(s)	Respirator ***
			In settings where caring for suspected/confirmed COVID-19 patient(s)	Well-fitting respirator or a medical mask
Known or suspected community or cluster transmission of SARS-CoV-2	Other staff, patients, visitors, service suppliers	Health care facility**	For any activity or in any common area	Well-fitting medical mask
	Inpatients		When physical distancing of at least 1 metre cannot be maintained or when outside of their care area	
	Health workers and caregivers	Health care facility**	In settings where caring for non-COVID-19 patients.	
		Home visit (for example, for antenatal or postnatal care, or for a chronic condition)		
Community	Community outreach programmes/essential routine services			
Known or suspected sporadic transmission of SARS-CoV-2 cases	Health workers and caregivers	Health care facility**	In settings when caring for non-COVID-19 patient(s)	Well-fitting medical mask
	Other staff, patients, visitors, service suppliers and all others		No routine activities in patient areas	A medical mask may not be required if no patient contact.

<b>No documented SARS-CoV-2 transmission</b>	Health workers and caregivers	Health care facility**	Providing any patient care	Medical mask use according to standard and transmission-based precautions
--	-------------------------------	------------------------	----------------------------	---

\*This table refers only to the use of medical masks and respirators. The use of medical masks and respirators should be combined with other personal protective equipment, standard and transmission-based precautions and other measures as appropriate, and always with hand hygiene.

\*\* Health facility can include primary, secondary, tertiary care levels, outpatient care, and long-term care facilities.

\*\*\* N95 or N99 or FFP2 or FFP3

Published 25 April 2022.

## Mask use in health care setting acknowledgements, authorship, contributions

### Funding

Funding for the development of this document was provided by the United States Centers for Disease Control and Prevention (US CDC), in addition to WHO core funds. However, the views expressed in this guidance do not necessarily reflect the official policies of the US CDC.

### Authorship, contributions, acknowledgements

WHO would like to thank the collaborative efforts of all those involved to make this process rapid, efficient, trustworthy and transparent.

#### *WHO Health Emergencies Programme COVID-19 Infection Prevention and Control Secretariat (alphabetically)*

**Benedetta Allegranzi** (WHO headquarters (WHO/HQ)), **April Baller** (WHO/HQ), **Kathy Dunn** (WHO/HQ), **Hannah Hamilton** (WHO/HQ), **Madison Moon** (WHO/HQ), **João Paulo Toledo** (WHO/HQ), **Victoria Willet** (WHO/HQ).

#### *WHO Health Emergencies Programme COVID-19 Infection Prevention and Control Steering Group (alphabetically)*

**Gertrude Avortri** (WHO Regional Office for Africa (AFRO)), **Deborah Barbasa** (AFRO), **Landry Cihambanya** (AFRO), **Astrid Lydia Chojnacki** (Regional Office for the Western Pacific (WPRO)), **Giorgio Cometto** (WHO/HQ), **Ana Paula Countinho Rehse** (Regional Office for Europe (EURO)), **Sergey Eremin** (WHO/HQ), **Ivan Ivanov** (WHO/HQ), **Maha Talaat Ismail** (Regional Office for the Eastern Mediterranean (EMRO)), **Luca Fontatna** (WHO/HQ), **Dennis Falzon** (WHO/HQ), **Nathan Ford** (WHO/HQ), **Pierre Claver Kariyo** (AFRO), **Catherine Kane** (WHO/HQ), **Mahmoud Hamounda** (EMRO), **Iman Heweidly** (EMRO), **Ying Ling Lin** (WHO/HQ), **Guy Mbayo** (AFRO), **Babacar Ndoye** (AFRO), **Bisso Hilde Fonse Okou** (AFRO), **Alice Simniceanu** (WHO/HQ), **Aparna Singh Shah** (Regional Office for South-Asia (SEARO)), **Nahoko Shindo** (WHO/HQ), **Howard Sobel** (WPRO), **Valeska Stempliuk** (Regional Office for the Americas/Pan American Health Organization (AMRO/PAHO)), **Maria Van Kerkhove** (WHO/HQ), **Puspha Ranhjan Wijesinghe** (SEARO), **Bassim Zayed** (EMRO), **Matteo Zignol** (WHO/HQ).

#### *UNICEF Observers*

**Nagwa Hasanin** (UNICEF), **Raoul Kamadjeu** (UNICEF), **Pierre Yves Oger** (UNICEF).

#### *WHO Health Emergencies Programme COVID-19 Infection Prevention and Control Guideline Development Group (alphabetically)*

**Yewanda Alimi** (Africa Centres for Disease Control and Prevention, Ethiopia), **Jameela Alsalman** (Ministry of Health of Bahrain, Bahrain), **Baba Aye** (Public Services International, France), **May C. Chu** (Colorado School of Public Health, Center for Global Health, United States of America), **John Conly** (Cumming School of Medicine, University of Calgary, Canada), **Barry David Cookson** (Division of Infection and Immunity, University College London, United Kingdom of Great Britain and Northern Ireland), **Nizam Damani** (Sindh Institute of Urology and Transplant Centre, Karachi, Pakistan (Dow University of Health Sciences, Pakistan), United Kingdom), **Dale Fisher** (Infectious Disease Division, Department of Medicine, National University Health System, Singapore), **Tiouri Benaissa Hanene** (Ministry of Health of Tunisia, Tunisia), **Joost Hopman** (Radboud University Medical Center, The Netherlands), **Mohammad**

**Mushtuq Husain** (Institute of Epidemiology Disease Control and Research (IEDCR), Bangladesh), **Kushlani Jayatilleke** (Sri Jayewardenepura General Hospital, Nugegoda, Sri Lanka, Sri Lanka), **Souha Kanj** (American University of Beirut Medical Center, Lebanon), **Daniele Lantagne** (Tufts University, United States of America), **Anna Levin** (University of São Paulo, Hospital das Clinicas, FM-USP, Brazil), **Yuguo Li** (Department of Mechanical Engineering, The University of Hong Kong, Hong Kong Special Administrative Region, China), **Moi Lin Ling** (Singapore General Hospital, SingHealth, Singapore), **Caline Mattar** (Division of Infectious Diseases, Washington University in St Louis (WUSTL), United States of America), **Mary-Louise McLaws** (Honorary clinical Epidemiology – University of New South Wales, Australia), **Geeta Mehta** (Journal of Patient Safety and Infection Control, India), **Shaheen Mehtar** (Infection Control Africa Network, South Africa), **Ziad Memish** (Ministry of Health of Saudi Arabia, Kingdom of Saudi Arabia), **Tochi Okwor** (Nigeria Centre for Disease Control, Nigeria), **Mauro Orsini** (Ministry of Health of Chile, Chile), **Diamantis Plachouras** (European Centre for Disease Prevention and Control (ECDC), Sweden), **Mathias W Pletz** (Institute for Infectious Diseases and Infection Control of the University Hospital of the Friedrich Schiller University, Germany), **Marina Salvadori** (Public Health Agency of Canada, Canada), **Ingrid Schoeman** (TB Proof, South Africa), **Mitchell Schwaber** (Ministry of Health of Israel, Israel), **Wing Hong Seto** (WHO Collaborating Centre for Infectious Disease Epidemiology and Control, School of Public Health, The University of Hong Kong, Hong Kong Special Administrative Region, China), **Mark Sobsey** (University of North Carolina and Aquagenx, LLC, United States of America), **Paul Ananth Tambyah** (National University of Singapore, Singapore), **Anucha Apisarntharak Thammsat** (University Hospital, Thailand), **Andreas Voss** (Radboud University Medical Centre, Nijmegen, NL, The Netherlands), **Walter Zingg** (Clinic for Infectious Diseases and Hospital Epidemiology, Zurich University Hospital, Zurich, Switzerland).

#### **Methodologist**

**Roger Chou** (Methodologist, Department of Medicine and Department of Medical Informatics and Clinical Epidemiology, Oregon Health & Science University, United States of America)

#### **Declaration of conflicts of interest**

Dr R. Chou is an author on some of the evidence used to inform some recommendations. However, as a methodologist, he provided guidance to the GDG on methodologic issues and is not a voting member of the GDG. In some meetings, he presented evidence and provided clarification on methods to guide discussions regarding the EtD tables; however, all decisions were made by voting members of the GDG.

#### **Temporary Advisor**

**Fernanda C. Lessa** (US Centers for Disease Control and Prevention, United States of America).

#### **Declaration of conflicts of interest**

Dr F. Lessa is an employee of the United States CDC which has provided funding towards the development of this guideline. After consulting with the WHO Ethics Committee, it was determined Dr Lessa would contribute to discussions as she brings significant technical and field expertise to the discussions; however, be recused from voting on recommendations.

#### **External reviewers (alphabetically)**

**Paul Hunter** (University of East Anglia, United Kingdom), **Kalisavar Marimuthu**, (National Centre for Infectious Diseases, Singapore Yong Loo Lin School of Medicine, National University of Singapore, Singapore), **Nalini Singh** (Department of Global Health, The Milken Institute School of Public Health, George Washington University)

**Declarations of interest of external reviewers were collected and assessed, and no conflict of interest was identified.**

## **PPE Technical Specifications**

Pertinent sections of the technical guidance, "[Technical specifications of personal protective equipment for COVID-19](#)", published 13 November 2020, will soon be incorporated in this living guidance.

**Technical specifications for medical masks [21]**

Item	Characteristics	Performance standards (or alternative equivalent)
Medical mask for a health care worker	Medical mask, good breathability, internal and external faces should be clearly identified, 98% droplet filtration, preferably fluid resistance.	Always use a clean, unused rectangular pleated medical mask meeting the following minimum performance standards (or equivalent): <ul style="list-style-type: none"> <li>• EN 14683 (Type II or Type IIR);</li> <li>• ASTM F2100 (Level 1, 2 or 3); or</li> <li>• YY 0469 OR YY/T 0969 (with at least 98% bacterial filtration efficiency).</li> </ul>
Medical mask for patient	Medical mask, good breathability, internal and external faces should be clearly identified	EN 14683 Type I YY 0469 or YY/T 0969, if bacterial droplet filtration is below 98% Or alternative equivalent standard

### Prevention, identification and management of health worker infection

The most up-to-date technical guidance for “Prevention, identification and management of health worker infection in the context of COVID-19” was published on 30 October 2020. This guidance is under review and is pending integration into [“Infection prevention and control in the context of coronavirus disease \(COVID-19\): A living guideline”](#).

### Rational use of PPE and considerations during severe shortages

The most up-to-date technical guidance for “Rational use of personal protective equipment for coronavirus disease (COVID-19) and considerations during severe shortages” was published on 23 December 2020. This guidance is under review and is pending integration into [“Infection prevention and control in the context of coronavirus disease \(COVID-19\): A living guideline”](#).

### Risk assessment and management of exposure

The most up-to-date technical guidance for “Risk assessment and management of exposure of health care workers in the context of COVID-19: interim guidance” was published 19 March 2020. This guidance is under review and is pending integration into [“Infection prevention and control in the context of coronavirus disease \(COVID-19\): A living guideline”](#).

### Safe dead body management

The most up-to-date guidance for [“Infection prevention and control for the safe management of a dead body in the context of COVID-19: interim guidance”](#) was published 4 September 2020. This guidance is under review and is pending integration into [“Infection prevention and control in the context of coronavirus disease \(COVID-19\): A living guideline”](#).

### Water, sanitation, hygiene, and waste management

The most up-to-date technical guidance for [“Water, sanitation, hygiene, and waste management for SARS-CoV-2, the virus that causes COVID-19”](#) was published 29 July 2020. This guidance is under review and is pending integration into [“Infection prevention and control](#)

[in the context of coronavirus disease \(COVID-19\): A living guideline](#)".

## Part 2: Community settings

Updated

Many of the existing technical guidance documents that will be integrated into this section are under review. Updated versions will be available in future versions. This section includes updated guidelines for mask use by the general public in community settings and mask use by children. Sections that are pending updates have links to the most recent iteration of relevant IPC guidance published online.

### Introduction to public health and social measures

#### What are PHSM?

PHSM have been implemented worldwide over the course of the pandemic to suppress SARS-CoV-2 transmission and reduce mortality and morbidity from COVID-19. PHSM include the following: personal protective measures (for example, physical distancing, avoiding crowded settings, hand hygiene, respiratory etiquette, mask-wearing); environmental measures (for example, cleaning, disinfection, ventilation); surveillance and response measures (for example, testing, genetic sequencing, contact tracing, isolation, and quarantine); physical distancing measures (for example, regulating the number and flow of people attending gatherings, maintaining distance in public or workplaces, domestic movement restrictions); and international travel-related measures. In this context, it does not include medical countermeasures such as drug administration or vaccination. PHSMs act in concert and a combination of measures is required to ensure adequate control. Measures should be implemented by the lowest administrative level for which situational assessment is possible and tailored to local settings and conditions. For more information, please refer to the [Considerations for implementing and adjusting public health and social measures in the context of COVID-19 \[65\]](#).

#### Adjusting PHSM

As the pandemic continues to evolve, PHSM should be regularly reviewed and adjusted according to the local epidemiology and its impact on the health system, including the community and the overall economy and society. This requires agile decision-making based on ongoing situational assessments at the most local administrative level possible in a coherent and coordinated manner with neighbouring areas at the sub-national and national levels. Such assessments should be based on available data and a risk/benefit approach considering the local epidemiology, the health system's capacity to respond and other contextual considerations (such as upcoming mass gathering events that may alter transmission or the health system's capacity). The choice of epidemiological indicators and their thresholds will depend on a country's data collection capacity, vaccination strategy and coverage, and the overall COVID-19 response strategy [65]. Important dynamic indicators to be considered to determine the local situation are SARS-CoV-2 transmissibility, the seriousness of COVID-19, and the impact on the health system. Assessments based on these key indicators (transmissibility, seriousness of disease, and impact) need to be tailored to the local context. As a general principle, core PHSM (for example, mask use, physical distancing) should be maintained in priority groups, settings and situations, even during periods of low transmission. By combining data regarding the above-mentioned three key indicators, the following situation levels can be identified to describe the local situation.

**Situational level 0:** A situation with no known transmission of SARS-CoV-2 in the preceding 28 days. The health system and public health authorities are ready to respond, but there are no restrictions needed on daily activities.

**Situational level 1:** A situation with minimal transmission, morbidity and health system impact of SARS-CoV-2, with only basic ongoing PHSM needed.

**Situational level 2:** A situation where there is a moderate impact of COVID-19, although there may be a higher impact in specific subpopulations. Additional measures may be required to reduce transmission. However, disruptions to social and economic activities can still be limited, particularly if PHSM can be targeted strategically to more impacted settings.

**Situational level 3:** A situation with a significant impact on the health system and a risk of health services becoming overwhelmed, or unacceptably high morbidity and mortality, despite sufficient remaining health system capacity. A broader combination of PHSM may need to be put in place to limit transmission, manage morbidity, and avoid overwhelming the health system.

**Situational level 4:** An uncontrolled epidemic with very high morbidity/mortality and limited or no additional health system response

capacity available, thus requiring extensive PHSM to avoid overwhelming of health services and substantial excess morbidity and mortality.

#### Who are these recommendations intended for?

These guidelines are intended for policy- and decision-makers, public health professionals, and IPC professionals at national, sub-national, and facility levels.

## Mask use

Updated

### Background

To assist national and global efforts to end the acute phase of the COVID-19 pandemic emergency worldwide, WHO published the 2022 COVID-19 Strategic Preparedness, Readiness and Response plan outlining strategic interventions to support these efforts. The first objective is to reduce and control the incidence of SARS-CoV-2 infections. This is essential to protect individuals from exposure, especially vulnerable individuals at risk of severe disease or occupational exposure to the virus, reduce the probability that future variants will arise, and reduce pressure on health systems. While the second objective is to prevent, diagnose and treat COVID-19 to reduce mortality, morbidity, and long-term sequelae [4]. These actions may reduce pressure on the virus to evolve and the potential that future variants will emerge while simultaneously reducing the burden on the health system [5].

Masks are one component of a comprehensive package of prevention and control measures to limit the spread of SARS-CoV-2. When aiming to reduce community transmission and mitigate the impact of COVID-19 outbreaks on health and social services, policies developed for mask use should be included as one element of a comprehensive package of preventive measures to reduce transmission (physical distancing, ventilation, mask use, hand hygiene, respiratory etiquette, and vaccination).

Considering the current stage of the pandemic, the GDG considered all available evidence on the effectiveness of mask-wearing [2], the epidemiology of current VoC, transmission (data or patterns where available), the severity of disease and impact on health systems, vaccine efficacy, access, uptake, and potential immune evasion [3]. The complementary strong and conditional recommendations on mask use in the community outline possible scenarios in which mask use may be of benefit.

## Mask use in the community

Strong recommendation for

Updated

Strong recommendation for, low to moderate certainty of evidence



WHO recommends the use of a mask for the prevention of SARS-CoV-2 transmission in the community in the following situations:

- when in crowded, enclosed, or poorly ventilated spaces<sup>1</sup> [16];
- following a recent exposure to COVID-19 (according to the WHO definition<sup>2</sup>) when sharing a space with others;
- when sharing a space with a person who displays signs or symptoms of COVID-19<sup>3</sup> or is COVID-19- positive;
- for individuals at high risk<sup>4</sup> of severe complications from COVID-19.

<sup>1</sup> For example, a setting in which it is not possible to physically distance at least 1 metre.

<sup>2</sup> Exposure: contact with a probable or confirmed case or linked to a COVID-19 cluster [66][67].

<sup>3</sup> Signs or symptoms of COVID-19 include: cough, general weakness/fatigue, headache, myalgia, sore throat, coryza, dyspnoea, nausea/diarrhoea/anorexia [66]

<sup>4</sup> High risk is defined as: people aged  $\geq 60$  years; or those with underlying comorbidities, such as cardiovascular disease or diabetes mellitus, chronic lung disease, cancer, cerebrovascular disease, immunosuppression, obesity, or asthma [68].

Published 13 January 2023

## Practical Info

### Implementation consideration

#### Implementation considerations

The implementation and adjustment of policies on mask use should be based on available scientific data and a risk/benefit approach considering the local epidemiology, the health system's capacity to respond, and other contextual considerations (events that may alter community transmission or the health system's capacity to respond to the resurgence of cases). The local situation can be determined based on the above-mentioned criteria related to the transmissibility of SARS-CoV-2, the seriousness of the disease, and the impact of the virus.

The evidence available on mask use in the community setting is based on the use of medical masks. Fabric (non-medical) masks can be used when access to medical masks is limited. While filtering facepiece respirators have demonstrated a higher filtration level, there is limited evidence to suggest that filtering facepiece respirators should be used in community settings.

Exhalation valves on respirators and non-medical masks are discouraged as they do not allow for adequate source control from the wearer. Exhalation valves permit a bypass of the filtration layers when the wearer exhales, thus potentially allowing infectious particles to pass through.

Face shields are considered to provide a level of eye protection only and should not be considered as an equivalent to masks with respect to respiratory protection and/or source control. Current laboratory testing standards only assess face shields for their ability to provide eye protection from chemical splashes [85].

#### Additional details

For additional information on the environmental impact of mask use (and other PPE), please see the WHO's Global analysis of health care waste in the context of COVID-19[106].

For information on assessing and improving indoor ventilation, please see WHO's Roadmap to improve and ensure good indoor ventilation in the context of COVID-19 [16].

For additional information on contact tracing and quarantine, please see Contact tracing and quarantine in the context of COVID-19: interim guidance, 6 July 2022 [67].

For the essential parameters concerning fabric (non-medical) and medical masks, see the following [implementation consideration](#).

## Evidence To Decision

### Benefits and harms

Substantial net benefits of the recommended alternative

The utilization of masks in community settings is associated with a decreased risk of SARS-CoV-2 infections compared with no mask-wearing. Despite the low-to-moderate certainty of the evidence, GDG members agreed that WHO should issue recommendations as the net benefits of mask use by the public outweigh the potential harms. The situations outlined above have been identified by consensus as settings and conditions in which masks should always be utilized.

Available evidence includes two open-label RCTs and ten observational studies. A large (n=342,183) cluster RCT found a mask promotion intervention associated with decreased risk of symptomatic SARS-CoV-2 seroprevalence (adjusted prevalence ratio 0.91, 95% CI 0.82 to 1.00) [69]. An individually randomized RCT (n=6,024) found a recommendation to use masks associated with decreased risk of SARS-CoV-2 infection, though the difference was not statistically significant (odds ratio 0.82, 95% CI 0.54 to 1.23); this trial was not designed to evaluate effects of masks as source control [70]. The RCTs had methodological limitations, including open-label design, attrition, incomplete outcome assessment, variable adherence, and differential recruitment. The RCTs were consistent and were not downgraded for imprecision (due to the very large total sample size [greatly exceeding any optimum information size threshold] with a precise estimate from one of the trials). Only one trial evaluated a mask recommendation directly [70]. The other evaluated a mask promotion intervention and did not evaluate mask use or a mask recommendation directly [69]; this resulted in suboptimal uptake of make use and would underestimate the effects of mask use. Therefore, the RCTs were not downgraded for indirectness (See Annex 2).

The observational studies were generally consistent with the RCTs, but had some imprecision, inconsistency and methodological limitations [71][72][73][74][75][76][77][78][79][80]. Although the estimates of the ten available observational studies were imprecise and had a degree of variability, in addition to other biases intrinsic to observational studies, overall, mask use was associated with a decreased risk of SARS-CoV-2 infection compared to no mask use [71][72][73][74][75][76][77][78][79][80]. Ecological studies identified an association between a reduced number of confirmed cases of COVID-19 and policies requiring the use of masks. No studies assessed the effectiveness of mask use in specific settings (for example, indoor, outdoor, or ventilation status). Overall, the certainty of the evidence (based primarily on the two RCTs, and supplemented by the ten observational studies) is assessed as *low-to-moderate*.

### Certainty of the Evidence

Available evidence includes two open-label RCTs and ten observational studies. The cluster RCT explored the use of mask promotion [69], while the other RCT presented an imprecise estimate and was not designed to assess the effectiveness of source control [70]. The observational studies had some imprecision, inconsistency and methodological limitations. Therefore, the certainty of the evidence is reported as *low-to-moderate*.

### Values and preferences

No substantial variability expected

Discussions with GDG members indicated a general preference towards favouring mask use in community settings, although the values and preferences of individuals may vary. Many members indicated that those at high risk of severe disease may find more value in the use of masks compared to other individuals.

**Resources**

No important issues with the recommended alternative

GDG members indicated that the global supply chain for mask manufacturing has improved and would not pose a severe obstacle to community masking. The cost of both medical masks and non-medical masks is relatively low and does not pose a substantial barrier for low- and middle-income countries. However, medical masks should not be reused and should be changed when wet or soiled, potentially requiring the use of multiple masks per day, leading to additional resource implications, such as cost, availability and access. Additionally, there are environmental impacts associated with disposable masks, such as additional waste and litter. Additional considerations are needed for proper disposal.

**Gaps in knowledge and research needs**

Investigations on the benefits and harms of masks and their utilization in the community setting are ongoing and published work has identified this need for continued research. Well-conducted, observational studies and/or RCTs exploring the use of masks versus no masks in various settings (for example, indoor, outdoor, ventilation status) would further clarify outstanding questions concerning mask use in community settings. In addition, research investigating the use of masks (including the type of mask and transmission scenarios) in the context of the emerging variants of concern would provide powerful evidence for future recommendations. However, GDG members discussed the challenges associated with obtaining compelling evidence from a RCT on behavioural interventions. Furthermore, with the availability of SARS-CoV-2 immunization and increased natural immunity, further research will be needed to reinforce the impact of vaccination and, consequently, the effect that immunization status will have on mask utilization in community settings. Additional research and innovation is needed in the area of reusable and recyclable medical masks that comply with existing standards.

**Equity**

No important issues with the recommended alternative

No issues were documented regarding inequities. Using masks as a preventative measure for SARS-CoV-2 infection may reduce the burden of infection, especially for those at high risk of severe disease [81]. Studies did not examine equity issues, such as providing information on race, gender, or vulnerable populations. More studies addressing these aspects should be carried out to inform the decision-making process.

**Acceptability**

Important issues, or potential issues not investigated

Complex issues arise when examining the acceptability of mask use in communities. These include the type of mask recommended, personal preference, possible local economic and procurement constraints and the ecological impact (environmental impact and waste management) [82][83][84]. Members indicated that those at high risk of severe sequelae might find more benefit in mask-wearing compared to other individuals. Furthermore, members of the general public may not deem mask use as an acceptable public health intervention and thus, demonstrate resistance towards masking policies. However, the evidence points to the benefits outweighing the harms. Variability exists in the published studies examining mask compliance.

**Feasibility**

Important issues, or potential issues not investigated

Given the current availability of masks, community masking is likely feasible, despite the potential acceptability issues mentioned above.

**Justification**

In response to the shift in the epidemiology of COVID-19, GDG members reformulated the recommendations to no longer rely on the local transmission scenario of SARS-CoV-2. Given the sustained SARS-CoV-2 transmission globally, a majority of GDG members agreed that a situational approach to mask use is more appropriate than the previous transmission-based approach. GDG members indicated that the benefits of mask use outweigh the potential harms as masks are an effective mitigation tool, especially in crowded, enclosed, and poorly ventilated settings such as public transportation, busy storefronts, and crowded workplaces and educational centres. The GDG decided for this strong recommendation in conjunction with the conditional to ensure coverage across all situations where masking may be beneficial.

Conditional recommendation for

Updated

### Conditional recommendation for, low to moderate certainty of evidence



In situations not addressed by the strong recommendation, WHO suggests a risk-based approach to inform the decision to use a mask for the prevention of SARS-CoV-2 transmission in the community.

Factors that favour mask use:

- COVID-19 epidemiological trends at the community level indicating high or rising transmission or hospitalizations;
- low coverage of COVID-19 vaccination;
- low levels of population immunity to SARS-CoV-2;
- a greater degree of crowding<sup>1</sup>, poorer indoor ventilation, and/or the presence of individual risk factors<sup>2</sup>.

<sup>1</sup> The degree of crowding for the conditional recommendation refers to distances >1 metre, for which there is likely some association between greater distancing and decreased risk.

<sup>2</sup> For the conditional recommendation, in the absence of clear risk factors, one may consider whether one's overall status of health may contribute to an increased risk of severe disease.

*Published 13 January 2023.*

## Practical Info

### Implementation consideration

#### Practical considerations for policy-makers:

The potential advantages of mask use by healthy people in the general public include:

- reduced spread of potentially infectious aerosols or droplets from exhaled breath, including from infected people before they develop symptoms [87];
- encouraging concurrent transmission prevention behaviours such as washing hands and not touching the eyes, nose and mouth [88][89][90]; and
- preventing transmission of other respiratory illnesses such as tuberculosis and influenza and reducing the burden of these diseases during the pandemic [91].

The potential disadvantages of mask use by healthy people in the general public include:

- Adverse events include: headache and/or breathing difficulties, depending on the type of mask used [92][93]; potential physiological changes [94]; development of facial skin lesions, irritant dermatitis or worsening acne when used frequently for long hours[93][95][96][97][98];
- difficulty with communicating clearly, especially for persons who are deaf or have poor hearing or use lip reading [99][100];
- poor compliance with mask-wearing, in particular by young children [95][101][102][103][104];
- waste management issues; improper mask disposal leading to increased litter in public places and environmental hazards [105][106]; and
- further disadvantages for, or difficulty wearing masks by, certain members of the population, especially: children; developmentally challenged people; those with mental illness or cognitive impairment; those with asthma, chronic

respiratory or breathing problems; those who have had facial trauma or recent oral maxillofacial surgery; and those living in hot and humid environments [92][95][102].

### Additional details

For additional information on the environmental impact of mask use (and other PPE), please see the WHO's Global analysis of health care waste in the context of COVID-19 [106].

For information on assessing and improving indoor ventilation, please see WHO's Roadmap to improve and ensure good indoor ventilation in the context of COVID-19 [16].

For additional information on contact tracing and quarantine, please see Contact tracing and quarantine in the context of COVID-19: interim guidance, 6 July 2022 [67].

For the essential parameters concerning fabric (non-medical) and medical masks, see the following [implementation consideration](#).

## Evidence To Decision

### Benefits and harms

Substantial net benefits of the recommended alternative

The utilization of masks in community settings is associated with a decreased risk of SARS-CoV-2 infections compared with no mask-wearing. Despite the low-to-moderate certainty of the evidence, GDG members agreed that WHO should issue recommendations as the net benefits of mask use by the public outweigh the potential harms. The situations outlined above have been identified by consensus as settings and conditions in which masks should always be utilized.

Available evidence includes two open-label RCTs and ten observational studies. A large (n=342,183) cluster RCT found a mask promotion intervention associated with decreased risk of symptomatic SARS-CoV-2 seroprevalence (adjusted prevalence ratio 0.91, 95% CI 0.82 to 1.00) [69]. An individually randomized RCT (n=6,024) found a recommendation to use masks associated with decreased risk of SARS-CoV-2 infection, though the difference was not statistically significant (odds ratio 0.82, 95% CI 0.54 to 1.23); this trial was not designed to evaluate effects of masks as source control [70]. The RCTs had methodological limitations, including open-label design, attrition, incomplete outcome assessment, variable adherence, and differential recruitment. The RCTs were consistent and were not downgraded for imprecision (due to the very large total sample size [greatly exceeding any optimum information size threshold] with a precise estimate from one of the trials). Only one trial evaluated a mask recommendation directly [70]. The other evaluated a mask promotion intervention and did not evaluate mask use or a mask recommendation directly [69]; this resulted in suboptimal uptake of make use and would underestimate the effects of mask use. Therefore, the RCTs were not downgraded for indirectness (See Annex 2).

The observational studies were generally consistent with the RCTs, but had some imprecision, inconsistency and methodological limitations [71][72][73][74][75][76][77][78][79][80]. Although the estimates of the ten available observational studies were imprecise and had a degree of variability, in addition to other biases intrinsic to observational studies, overall, mask use was associated with a decreased risk of SARS-CoV-2 infection compared to no mask use [71][72][73][74][75][76][77][78][79][80]. Ecological studies identified an association between a reduced number of confirmed cases of COVID-19 and policies requiring the use of masks. No studies assessed the effectiveness of mask use in specific settings (for example, indoor, outdoor, or ventilation status). Overall, the certainty of the evidence (based primarily on the two RCTs, and supplemented by the ten observational studies) is assessed as *low-to-moderate*.

### Certainty of the Evidence

Available evidence includes two open-label RCTs and ten observational studies. The cluster RCT explored the use of mask promotion [69], while the other RCT presented an imprecise estimate and was not designed to assess the effectiveness of source control [70]. The observational studies had some imprecision, inconsistency and methodological limitations. Therefore, the certainty of the evidence is reported as *low-to-moderate*.

**Values and preferences**

Substantial variability is expected or uncertain

Discussions with GDG members indicated a general preference towards favouring mask use in community settings, although the values and preferences of individuals may vary. Many members indicated that those at high risk of severe disease might perceive the benefits of mask use to be greater compared to other individuals.

**Resources**

No important issues with the recommended alternative

Many GDG members noted that the global supply chain for mask manufacturing has improved and would not pose a severe obstacle to community masking. The cost of both medical and non-medical (fabric) masks is relatively low and does not pose a substantial barrier for low- and middle-income countries. However, medical masks should not be reused and changed when wet or soiled, potentially requiring the use of multiple masks per day, leading to additional resource implications, such as cost, availability and access. Additionally, there are environmental impacts associated with disposable masks, such as additional waste and litter. Additional considerations are needed for proper disposal.

**Gaps in knowledge and research needs**

Investigations on the utilization of masks in the community setting are ongoing, but published work has identified this need for continued research. Observational studies and/or RCTs designed and conducted with rigorous scientific methods exploring the use of masks versus no masks in various settings (for example, indoor, outdoor, ventilation status) would further clarify outstanding questions concerning mask use in community settings. In addition, research investigating the use of masks (including the type of mask and transmission scenarios) in the context of VoC would provide powerful evidence for future recommendations. However, GDG members discussed the challenges associated with obtaining compelling evidence from an RCT on behavioural interventions. Furthermore, with the availability of SARS-CoV-2 immunization and increases in natural immunity, further research will be needed to reinforce the impact of vaccination and; consequently, the effect immunization status will have on mask utilization in community settings. Additional research and innovation is needed in the area of reusable and recyclable medical masks that comply with existing standards.

**Equity**

No important issues with the recommended alternative

No important issues were documented regarding inequities, although this arena would benefit from further investigation.

**Acceptability**

Important issues, or potential issues not investigated

Complex issues arise when examining the acceptability of mask use in communities. These include the type of mask recommended, personal preference and ecological impact (environmental impact and waste management) [82][83][84][106]. Members indicated that those at high risk of severe sequelae might find more benefit in mask-wearing compared to other individuals. Furthermore, it has been indicated that members of the general public may not deem mask use as an acceptable public health intervention and, thus, demonstrate resistance towards masking policies. However, the evidence points to the benefits outweighing the harms. Variability exists in the published studies examining mask compliance.

**Feasibility**

Important issues, or potential issues not investigated

Given the availability of masks, community masking is likely feasible.

**Justification**

GDG members decided for this conditional recommendation in conjunction with the aforementioned strong recommendation. In addition to situations where masks are strongly advised (when in crowded, enclosed, or poorly ventilated spaces; following recent exposure to COVID-19; when sharing a space with a person who displays symptoms of COVID-19 or is COVID-19-positive; and for individuals at high risk of severe complications from COVID-19), there are additional times where wearing a mask may be

beneficial. Although there are limited data on the effectiveness of a risk-based approach and implementation may be a challenge, the benefits of mask wearing outweigh the risks.

Members indicated that masks should be considered when there are high-to-moderate levels of community transmission (situational levels 2 to 4) and low-to-moderate vaccination coverage while taking into consideration individual risk factors in addition to personal values and preferences based on the perception of the risk and the potential harm and consequences of being affected by COVID-19.

#### Implementation consideration

##### Mask management

For any type of mask, appropriate use, storage, cleaning or disposal are essential to ensure that they are as effective as possible and to avoid any increased risk of transmission. Adherence to correct mask management practices varies, reinforcing the need for appropriate messaging [86]. WHO provides the following guidance on the correct use of masks:

- Wash hands thoroughly before putting on the mask.
- Inspect the mask for tears or holes, and do not use a damaged mask.
- Place the mask carefully, ensuring it covers the mouth and nose, adjust to the nose bridge and tie it securely to minimize any gaps between the face and the mask. If using ear loops, ensure these do not cross over as this widens the gap between the face and the mask.
- Avoid touching the mask while wearing it. If the mask is accidentally touched, wash hands thoroughly.
- Remove the mask using the appropriate technique. Do not touch the front of the mask; rather, untie it from behind.
- Replace the mask as soon as it becomes damp with a new, clean and dry mask.
- Either discard the mask or place it in a clean plastic resealable bag where it is kept until it can be washed and cleaned. Do not store the mask around the arm or wrist or pull it down to rest around the chin or neck.
- Wash hands immediately after discarding a mask.
- Do not reuse single-use masks.
- Discard single-use masks after each use and properly dispose of them immediately upon removal.
- Do not remove the mask to speak.
- Do not share your mask with others.
- Wash fabric masks in soap or detergent and preferably hot water (at least 60° Centigrade/140° Fahrenheit) at least once a day. If it is not possible to wash the masks in hot water, then wash the mask in soap/detergent and room-temperature water, followed by boiling the mask for 1 minute.
- A mask should be changed to a clean mask at least once daily.

For more information on mask technical specifications, review the following technical document - "[Technical specifications of personal protective equipment for COVID-19](#)", published 13 November 2020

---

*Published 13 January 2023.*

Good practice statement

Updated



Individuals with any signs or symptoms<sup>1</sup> suggestive of COVID-19 or who test positive for COVID-19 should wear a medical mask, when sharing a space with others, until it is resolved or the isolation period is complete.

<sup>1</sup> Signs or symptoms of COVID-19 include: cough, general weakness/fatigue, headache, myalgia, sore throat, coryza, dyspnoea, nausea/diarrhoea/anorexia [66].

Published 13 January 2023.

## Practical Info

### Implementation consideration

- Individuals should self-isolate and seek medical advice as soon as they start to feel unwell with potential COVID-19 symptoms (even if symptoms are mild).
- Instructions on how to put on, take off and dispose of medical masks, and how to adequately perform hand hygiene [107] should be followed.
- All additional measures should be followed, particularly respiratory hygiene, frequent hand hygiene, and maintaining a physical distance of at least one metre from other persons [108].
- If a medical mask is not available for individuals with suspected or confirmed COVID-19, a fabric mask with fit, filtration and breathability assessed to meet WHO's essential parameters for non-medical masks should be worn by patients as a source control measure, pending access to a medical mask. The use of a non-medical mask can minimize the projection of respiratory particles from the user [109][110].
- Persons with suspected COVID-19 or mild COVID-19 symptoms should wear a medical mask as much as possible, especially when there is no alternative to being in the same room with other people.
- Caregivers or those sharing living space with people with suspected COVID-19 or mild COVID-19 symptoms should wear a medical mask when in the same room as the affected person.

## Justification

GDG members agreed that if an individual has confirmed or suspected COVID-19 needs to interact with others in or outside of their household, they should wear a medical mask. Members also noted that individuals who have confirmed or suspected COVID-19 should self-isolate for the duration of their isolation period and/or until symptoms resolve. For additional information on contact tracing and quarantine, please see Contact tracing and quarantine in the context of COVID-19: interim guidance, 6 July 2022 (who.int) [67].

Good practice statement

Updated



Policies aimed at reducing the transmission of SARS-CoV-2 in the community should be revisited, strengthened, and updated according to the most recent scientific evidence.

Published 13 January 2023.

## Practical Info

Policies may include a package of interventions such as vaccination, ventilation, physical distance, hand hygiene, respiratory etiquette, and mask adherence by the general public. Please refer to the document on IPC in the event of surge or resurgence in cases of COVID-19 [167].

## Justification

GDG members agreed that national and subnational policy-makers should revisit, strengthen and update local policies according to the most recent scientific evidence to mitigate SARS-CoV-2 transmission in the community settings. These policies should be able to be quickly scaled up should COVID-19 incidence increase in the community and if healthcare systems are at risk of becoming overwhelmed. Policies should be reviewed as necessary to account for any changes in the local context or new VoC.

## Type of mask used by the general public

Implementation consideration for policy-makers, when providing guidance, or setting standards for manufacturers on type of mask used by the general public

Updated

### Implementation consideration

The following mask types are acceptable options for use by the general public:

- disposable medical masks, if the availability of medical masks meeting minimum performance criteria for health workers has been assured\*;
- non-medical masks that comply with standards for safety and efficacy\*\* and can be washed prior to reuse;
- if the above options are not available, other types of well-fitting non-medical masks\*\*\* are an acceptable option (according to local policies).

\*Complying with medical mask standards (at minimum) EN 14683 type I, ASTM F2100 level 1, YY/T 0969, YY 0469 (or equivalent). For requirements for health workers, please see PPE technical specifications.

\*\*Complying with the ASTM F3502-22a Standard Specification Specification for Barrier Face Coverings, standard or a non-medical mask meeting WHO essential parameters (see *Practical information* for more information).

\*\*\*Including homemade, multi-layered masks (see *Practical information* for more information).

Published 13 January 2023.

## Practical Info

Table 2. Essential parameters (minimum and preferred thresholds) for manufactured non-medical mask

Essential Parameters	Minimum threshold	Preferred threshold
<b>1. Filtration*</b>		
<b>1.1 Filtration efficiency</b>	70% at 3 µm	>50% at 0.3 µm, without compromising breathability
<b>1.2. Challenge particle</b>	<i>Solid:</i> sodium chloride (NaCl), Talcum powder, Holi powder, dolomite, Polystyrene Latex spheres <i>Liquid:</i> DEHS Di-Ethyl-Hexyl-Sebacat, paraffin oil	<i>Solid:</i> sodium chloride (NaCl), Polystyrene Latex spheres
<b>1.3. Particle size</b>	Choose either size: 3 µm, 1 µm, or smaller	0.3 µm
<b>2. Breathability</b>		

<b>2.1. Breathing resistance**</b>	$\leq 70 \text{ Pa/cm}^2$	<p><i>Adult: <math>\leq 40 \text{ Pa/cm}^2</math></i></p> <p><i>Children: <math>\leq 20 \text{ Pa/cm}^2</math></i></p>
<b>2.2 Exhalation valves</b>	Not recommended	N/A
<b>3. Fit</b>		
<b>3.1. Coverage</b>	Full coverage of nose and mouth, consistent, snug perimeter fit at the nose bridge, cheeks, chin and lateral sides of the face; adequate surface area to minimize breathing resistance and minimize side leakage	Same as current requirements
<b>3.2 Face seal</b>	Not currently required	<p>Seal as good as FFR (respirator)</p> <p>Fit factor of 100 for N95</p> <p>Maximum Total Inward Leakage of 25% (FFP1 requirement)</p> <p>OR</p> <p>Leakage ratio of <math>\geq 5</math></p>
<b>3.2. Sizing</b>	Adult and child	<p>Should cover from nose bridge to below the chin and cheeks on either side of the mouth</p> <p>Sizing for adults and children (6-9, 10-12, &gt;12)</p>
<b>3.3 Strap strength</b>		> 44.5 N

\* Smaller particles may result in lower filtration.

\*\* High resistance can cause bypass of the filtration layers of the mask. Unfiltered air will leak out the sides or around the nose on the path of least resistance.

**Table 3. Additional (optional) parameters for manufactured non-medical masks**

<b>Additional parameters</b>	<b>Minimum thresholds</b>
If reusable, the number of wash cycles	5 cycles
Disposal	If majority of mask is compostable, as per EN 13432, EN 14995, ASTM D5511 or other similar standards mimicking landfill or marine environments
Antimicrobial (bacteria, virus, fungus) performance	<p>ISO 18184 (virus)</p> <p>ISO 20743 (bacteria)</p> <p>ISO 13629 (fungus)</p>

	AATCC TM100 (bacteria)
Chemical safety	Comply with REACH regulation, including inhalation safety

**Standards organizations' performance criteria**

Manufacturers producing masks with consistent standardized performance can adhere to published, freely available guidance from several organizations including those from, ASTM International, the French Standardization Association (AFNOR Group), The European Committee for Standardization (CEN), Swiss National COVID-19 Task Force, the South Korean Ministry of Food and Drug Safety (MFDS), the Italian Standardization Body (UNI) and the Bangladesh Directorate General of Drug Administration (DGDA).

**Additional criteria:**

- The non-medical mask, including all components and packaging, must be non-hazardous, non-toxic and child-friendly (no exposed sharp edges, protruding hardware or rough materials).
- Factory-made EN Type I, ASTM Level 1 medical masks or non-medical masks must be made using a process that is certified to a quality management system (e.g., ISO 13485, ISO 9001).
- Social accountability standards (e.g., SAI SA8000) for multiple aspects of fair labour practices, health and safety of the workforce and adherence to UNICEF's Children's Rights and Business Principles are strongly encouraged.




Figure 1. Illustration of the three essential parameters of filtration, breathability and fit.

*Filtration and breathability*

Filtration depends on the filtration efficiency (in %), the type of challenge particle (oils, solids, droplets containing bacteria) and the particle size (see Table 2). Depending on the fabrics used, filtration and breathability can complement or work against each other. Filtration is dependent on the tightness of the weave, fibre or thread diameter. Non-woven materials used for disposable masks are manufactured using processes to create polymer fibres that are thinner than natural fibres such as cotton, and that are held together by partial melting. Breathability is the difference in pressure across the mask and is typically reported in millibars (mbar) or Pascals (Pa) or normalized to the cm<sup>2</sup> in mbar/cm<sup>2</sup> or Pa/cm<sup>2</sup>. Non- medical fabric masks consisting of two layers of polypropylene spunbond, and two layers of cotton have been shown to meet the minimum requirements for droplet filtration and breathability of the CEN/TS 17553:2022 guidance. It is preferable not to select elastic material to make masks as the mask material may be stretched over the face, resulting in increased pore size and lower filtration through reuse. Additionally, elastic fabrics are sensitive to washing at high temperatures and may therefore degrade over time.

Coating the fabric with compounds such as wax may increase the barrier and render the mask fluid-resistant; however, such coatings may inadvertently block the pores completely and make the mask difficult to breathe through. In addition to decreased breathability, unfiltered air may more likely escape from the sides of the mask on exhalation. The coating is

therefore not recommended.

#### *Fit: shape and sizing*

Fit is the third essential parameter, and takes into consideration coverage, seal, sizing and strap strength. Fit of masks is currently not defined by any standard except for the anthropometric considerations of facial dimensions (ISO/TS 16976-2) or simplified to height mask (South Korean standard for KF-AD). Ideally, the mask should not have contact with the lips, unless hydrophobic fabrics are used in at least one layer of the mask [168]. Leaks where unfiltered air moves in and out of the mask may be attributed to the size and shape of the mask [169].

#### **Optional parameters for consideration**

If reusable:

- the biodegradability;
- antimicrobial performance (where applicable); and
- chemical safety (see Practical Info section).

Non-medical masks intended to be reusable should include instructions for washing and must be washed a minimum of five cycles, implying initial performance is maintained after each wash cycle. Advanced fabrics may be biodegradable or compostable at the end of service life, according to a recognized standard process (e.g., UNI EN 13432, UNI EN 14995 and UNI/PdR 79).

Manufacturers sometimes claim their non-medical masks have antimicrobial performance. Antimicrobial performance may be the result of coatings or additives to the fabric fibres. Treated fabrics must not come into direct contact with mucous membranes; the innermost fabric should not be treated with antimicrobial additives, only the outermost layer. In addition, antimicrobial fabric standards (e.g., ISO 18184, ISO 20743, AATCC TM100, AATCC 100) are generally slow acting. The inhibition on microbial growth may not take full effect until after a contact time of 2–24 hours, depending on the standard. The standards have generally been used for athletic apparel and to substantiate claims of odour control performance. These standards are not appropriate for non-medical cloth masks and may provide a false sense of protection from infectious agents. If claims are made, manufacturers should specify the standard that supports antimicrobial performance, the challenge organism and the contact time.

Volatile additives are discouraged as these may pose a health risk when inhaled repeatedly during wear. Certification according to organizations including OEKO-TEX (Europe) or SEK (Japan), and additives complying with REACH (Europe) or the United States Environmental Protection Agency (EPA), indicate that textile additives are safe and added at safe levels.

#### **Justification**

GDG members agreed with standardizing recommendations for the utilization and specifications of masks for the general public. GDG members expressed concern of being overly prescriptive while the current state of evidence on the quality and effectiveness of non-medical masks continues to evolve, as this may limit the social enterprise of homemade mask production, a standard practice within many WHO Member States. However, GDG members agreed with laboratory evidence confirming that non-medical masks without standardized quality control processes can have large variabilities in their key parameters (see *practical information* for information on essential parameters for non-medical masks). Members also conveyed the importance of specifying the use of well-fitting masks, as the fit may be an essential parameter for effective source control and protection. In addition, GDG members discussed the potential harms associated with limited resources and lack of personnel to test the essential parameters of masks in various low-income settings, together with expressing concerns regarding waste disposal.

#### **Adaptation**

**Homemade non-medical masks made from household fabrics (e.g. cotton, cotton blends and polyesters) should ideally have a three-layer structure, with each layer providing a function (see Figure 1) [83].**

1. an innermost layer (that will be in contact with the face) of a hydrophilic material (e.g. cotton or cotton blends of terry cloth towel, quilting cotton and flannel) that is non-irritating against the skin and can contain droplets [168];

2. a middle hydrophobic layer of synthetic breathable non-woven material (spunbond polypropylene, polyester and polyaramid), which may enhance filtration, prevent permeation of droplets or retain droplets [168][84]; and
3. an outermost layer made of hydrophobic material (e.g. spunbond polypropylene, polyester or their blends), which may limit external contamination from penetrating through the layers to the wearer's nose and mouth and maintains and prevents water accumulation from blocking the pores of the fabric [168].




Figure 1. Non-medical mask construction using breathable fabrics such as cotton, cotton blends, polyesters, nylon and polypropylene spunbond that are breathable may impart adequate filtration performance when layered. Single- or double-layer combinations of advanced materials may be used if they meet performance requirements [85]



Although a minimum of three layers is recommended for non-medical masks for the most common fabric used, single, double or other layered combinations of advanced materials may be used if they meet performance requirements.

Assumptions regarding homemade masks are that individual makers only have access to common household fabrics and do not have access to test equipment to confirm target performance (filtration and breathability). Figure 1 illustrates a multi-layer mask construction with examples of fabric options. Very porous materials, such as gauze, even with multiple layers, may provide very low filtration efficiency [86]. Fabrics with higher thread count offer improved filtration performance [87]. Coffee filters, vacuum bags and materials not meant for clothing should be avoided, as they may contain injurious content when breathed in. Microporous films such as Gore-Tex are not recommended [88].

## Mask use during physical activity


In review

WHO advises that people should not wear masks during vigorous-intensity physical activity [111] because masks may reduce the ability to breathe comfortably. The most important preventive measure is to maintain physical distancing of at least 1 metre and to ensure good ventilation when exercising.

*Extracted from the guidance titled "Mask use in the context of COVID-19", published 1 December 2020.*

## Practical Info

When community or cluster transmission of SARS-CoV-2 is experienced in local context, particular attention should be paid to ensuring physical distancing of at least 1 metre between persons outside of their households and frequent cleaning and disinfection of any public environment in which exercise is performed, especially high-touch surfaces. As well, if the activity takes place indoors, adequate ventilation (e.g. 10 litres of air exchange per second, per person occupying an indoor space) should be ensured at all times through natural ventilation or a properly functioning and maintained ventilation system [115]. If all the above measures cannot be ensured, consider temporary closure of public indoor exercise facilities (e.g. gyms).



## Evidence To Decision

### Benefits and harms

There are limited studies on the benefits and harms of wearing medical masks, respirators and non-medical masks while exercising. Several studies have demonstrated statistically significant deleterious effects on various cardiopulmonary physiologic parameters during mild to moderate exercise in healthy subjects and in those with underlying respiratory diseases [112][113][114][116][117][118]. The most significant impacts have been consistently associated with the use of respirators and in people with underlying obstructive airway pulmonary diseases such as asthma and chronic obstructive pulmonary disease (COPD), especially when the condition is moderate to severe [114]. Facial microclimate changes with increased temperature, humidity and perceptions of dyspnoea were also reported in some studies on the use of masks during exercise [113][119]. A recent review found negligible evidence of any negative effects of mask use during exercise but noted concern for individuals with severe cardiopulmonary disease [120].

## Mask use in the community authorship, acknowledgements, and contributors

### **Authorship, contributions and acknowledgments**

WHO would like to thank the collaborative efforts of all those involved to make this process rapid, efficient, trustworthy and transparent.

### **WHO Health Emergencies Programme COVID-19 Infection Prevention and Control Secretariat (alphabetically)**

**Benedetta Allegranzi** (WHO/HQ), **April Baller** (WHO/HQ), **Kathy Dunn** (WHO/HQ), **Nathan Ford** (WHO/HQ), **Hannah Hamilton** (WHO/HQ), **João Paulo Toledo** (WHO/HQ).

### **WHO Health Emergencies Programme COVID-19 Infection Prevention and Control Steering Group (alphabetically)**

**Lisa Askie** (WHO/HQ), **Anshu Banerjee** (WHO/HQ), **Adriana Velazquez Berumen** (WHO/HQ), **Astrid Lydia Chojnacki** (WPRO), **Landry Kabego Cihambanya** (AFRO), **Jennifer Collins** (EURO), **Giorgio Cometto** (WHO/HQ), **Janet Victoria Diaz** (WHO/HQ), **Sergey Eremin** (WHO/HQ), **Dennis Falzon** (WHO/HQ), **Luca Fontana** (WHO/HQ), **Melinda Frost** (WHO/HQ), **Bruce Allan Gordon** (WHO/HQ), **Iman Heweidy** (EMRO), **Maha Talaat Ismail** (EMRO), **Ivan Dimov Ivanov** (WHO/HQ), **Kathryn Johnston** (PAHO), **Ying Ling Lin** (WHO/HQ), **Tendai Makamure** (AFRO), **Madison Moon** (WHO/HQ), **Leandro Pecchia** (WHO/HQ), **Mark Perkins** (WHO/HQ), **Ana Paula Coutinho Rehse** (EURO), **Nahoko Shindo** (WHO/HQ), **Alice Simniceanu** (WHO/HQ), **Aparna Singh Shah** (SEARO), **Valeska Stempliuk** (PAHO), **Victoria Willet** (WHO/HQ), **Maria Van Kerkhove** (WHO/HQ).

### **UNICEF Observers**

**Nagwa Hasanin** (UNICEF), **Raoul Kamadjeu** (UNICEF), **Pierre Yves Oger** (UNICEF).

### **WHO Health Emergencies Programme COVID-19 Infection Prevention and Control Guideline Development Group (alphabetically)**

**Yewanda Alimi** (Africa Centres for Disease Control and Prevention, Ethiopia), **Jameela Alsalman** (Ministry of Health of Bahrain, Bahrain), **Baba Aye** (Public Services International, France), **May C. Chu** (Colorado School of Public Health, Center for Global Health, United States of America), **John Conly** (Cumming School of Medicine, University of Calgary, Canada), **Barry David Cookson** (Division of Infection and Immunity, University College London, United Kingdom of Great Britain and Northern Ireland), **Nizam Damani** (Sindh Institute of Urology and Transplant Centre, Karachi, Pakistan (Dow University of Health Sciences, Pakistan), United Kingdom), **Fernanda C. Lessa** (US Centers for Disease Control and Prevention, United States of America), **Dale Fisher** (Infectious Disease Division, Department of Medicine, National University Health System, Singapore), **Tiouiri Benaissa Hanene** (Ministry of Health of Tunisia, Tunisia), **Mohammad Mushtuq Husain** (Institute of Epidemiology Disease Control and Research (IEDCR), Bangladesh), **Kushlani Jayatilleke** (Sri Jayewardenepura General Hospital, Nugegoda, Sri Lanka, Sri Lanka), **Souha Kanj** (American University of Beirut Medical Center, Lebanon), **Daniele Lantagne** (Tufts University, United States of America), **Anna Levin** (University of São Paulo, Hospital das Clinicas, FM-USP, Brazil), **Yuguo Li** (Department of Mechanical Engineering, The University of Hong Kong, Hong Kong Special Administrative Region, China), **Moi Lin Ling** (Singapore General Hospital, SingHealth, Singapore), **Caline Mattar** (Division of Infectious Diseases, Washington University in St Louis (WUSTL), United States of America), **Mary-Louise McLaws** (Honorary clinical Epidemiology – University of New South Wales, Australia), **Geeta Mehta** (Journal of Patient Safety and Infection Control, India), **Shaheen Mehtar** (Infection Control Africa Network, South Africa), **Ziad Memish** (Ministry of Health of Saudi Arabia, Kingdom of Saudi Arabia), **Tochi Okwor** (Nigeria Centre for Disease Control, Nigeria), **Diamantis Plachouras** (European Centre for Disease Prevention and Control (ECDC), Sweden), **Mathias W Pletz** (Institute for Infectious Diseases and Infection Control of the University Hospital of the Friedrich Schiller University, Germany), **Marina Salvadori** (Public Health Agency of Canada, Canada), **Ingrid Schoeman** (TB Proof, South Africa), **Mitchell Schwaber** (Ministry of Health of Israel, Israel), **Mark Sobsey** (University of North Carolina and Aquagenx, LLC, United States of America), **Paul Ananth Tambyah** (National University of Singapore, Singapore), **Walter Zingg** (Clinic for Infectious Diseases and Hospital Epidemiology, Zurich University Hospital, Zurich, Switzerland).

### **Methodologist**

**Roger Chou** (Methodologist, Department of Medicine and Department of Medical Informatics and Clinical Epidemiology, Oregon Health & Science University, United States of America)

### **Declaration of conflicts of interest**

*Dr R. Chou* is an author on some of the evidence used to inform some recommendations. However, as a methodologist, he provided guidance to the GDG on methodologic issues and is not a voting member of the GDG. In some meetings, he presented evidence and provided clarification on methods to guide discussions regarding the EtD tables; however, all decisions were made by voting members of the GDG.

### **External reviewers (alphabetically)**

**Paul Hunter** (University of East Anglia, United Kingdom), **Mark Loeb** (McMaster University, Canada), **Nalini Singh** (Department of Global Health, The Milken Institute School of Public Health, George Washington University)

*Declarations of interest of external reviewers were collected and assessed, and no conflict of interest was identified.*

## Mask use by children

### Guiding Principles

Given the limited evidence on the use of masks by children in the context of COVID-19, including limited evidence on transmission of SARS-CoV-2 in children at specific ages, policy formulation by national authorities should be guided by the following overarching principles.:

- Do no harm: the best interest, health and well-being of the child should be prioritized.
- The application of these guidelines should not impact development or learning outcomes, including access to education.
- The guidelines should consider the feasibility of implementing recommendations in different social, cultural and geographic contexts, including limited resource and humanitarian settings, and among children with disabilities or specific health conditions.
- Any recommendation for mask use for children should encompass needed flexibility to enable children to maintain their rights to play, to education and ability to engage in everyday activities [19].
- National policies on the use of masks for children should be adapted based on social, cultural and environmental considerations, including in settings with limited resources and humanitarian settings.

## Introduction

## Introduction

WHO guidance on the use of masks for children in the community was first published in August 2020 as an annex to the document [Mask use in the context of COVID-19 \[7\]\[9\]](#). In December 2021, it was incorporated into the online version 1.0 of the WHO IPC COVID-19 living guideline published using the MAGICapp platform [6]. This updated version includes new recommendations for mask use by children of different ages, accommodations for children living with disabilities and updated implementation considerations, including for school settings.

WHO and UNICEF jointly developed this guideline. A guideline development group, the *WHO-UNICEF GDG for the use of masks by children in the context of COVID-19*, was established. Details on the composition of the GDG and the retrieval, synthesis and assessment of evidence can be found in the [methods](#) and [acknowledgements](#) sections of the document.

When aiming to reduce community transmission and mitigate the impact of COVID-19 outbreaks on health and social services, policies developed for mask use should be included as one element of a comprehensive package of preventive measures to reduce transmission (ventilation, physical distance, hand hygiene, and respiratory etiquette). In any decision being made related to the use of masks by children, the guiding principles for the best interest of children and a “do no harm” approach should prevail.

Each country is facing a different situation in the pandemic depending on a number of factors including the intensity of SARS-CoV-2 circulation, amount of population level immunity, capacities to respond and agility to adjust measures. As the pandemic continues and the virus evolves, changes in transmission intensity, the circulating variant of concern, and the capacities for health systems to respond based on the situation will result in need for policy adjustments related to IPC and PHSM. National policies should be evidence based, agile and adjusted as needed taking into consideration these and other factors. Countries should conduct an assessment of the transmission scenario and the health system response capacity – and assign a situational level to a geographic area. The assessment should examine quantitative and qualitative information from available sources, and can refer to the situational and community transmission (CT) Levels CT1-CT4 as described in, [Considerations for implementing and adjusting public health and social measures in the context of COVID-19 \[65\]](#). Additional factors, including population level immunity, will need to be taken into account when setting national and sub-national policies, as outlined above.

This section of the guideline focuses on the use of masks in children in the context of COVID-19 in community settings, such as schools and recreational areas. Children spend a considerable portion of their time in schools, which may have indoor and outdoor areas, and there are existing specific guidance documents available that address school-related public health measures.

Recommendations on types of masks can be found in the [mask use in community settings](#) section of the document.

There are five statements for the use of masks by children, including three recommendations by [age group](#) ([≤ 5](#), [6-11](#), [12 and over](#)), and [two good practice statements](#).

## Evaluation

### Monitoring and evaluation of the impact of mask use by children

When implementing policies for mask-wearing for children, key information should be collected on a regular basis and where possible utilized to inform future policy. Monitoring and evaluation should be established at the onset and include:

- indicators that measure the impact on the child’s health, including mental health
- reduction in transmission of SARS-CoV-2 at community and health facility level
- motivators and barriers to mask-wearing
- impacts on children’s development and learning and school attendance
- ability of children to express themselves
- impact on children with developmental delays, health conditions and disabilities or other vulnerabilities
- experiences of children, their needs, perspectives and expectations.

Data should be used to inform policy updates and strategies on:

- communication

- training and support to teachers, educators, parents and children
- distribution of materials that empower children to use masks appropriately
- indicators to lift mask requirements for children.

Analyses should include sex, age and physical, social and economic stratification to ensure that policy implementation reduces health and social inequities.

### Research Needs

There are significant limitations in the available evidence on benefits and harms of mask use in children including a lack of evidence on important developmental and long-term outcomes. Future studies should consider evaluation of the effectiveness of mask use by children of different age groups in reducing transmission of SARS-CoV-2, impacts on learning and development, psychological health and quality of life. While RCTs would be ideal, well conducted observational studies that control for other infection control measures, exposures and other confounders would also be informative.

#### SARS-CoV-2 Transmission in Children

Disease severity and mortality due to COVID-19 including infections with VOCs increases with age, and children tend to present with a milder course of illness than older population groups [122][3][123]. The transmission characteristics among children need to be interpreted in light of new VOC's, in particular, Omicron; vaccination strategies and age-specific vaccination coverage and changes in mixing patterns as a result of the implementation of PHSM. Evidence early in the pandemic from household, serological and infection prevalence studies suggested that young children may be at lower risk of infection than adolescents and adults and potentially transmit SARS-CoV-2 less [122][123][124][125][126][127][128][129][130][131][132][133][134]. However, more recent epidemiological trends seem to indicate that children contribute to transmission similarly to adults, due to their social mixing patterns in some settings and in light of emerging VOC's such as Omicron [42][135][136][137][138][139]. This has been well documented in settings where extensive community testing has been undertaken (e.g. the REACT study in the United Kingdom) [140]. The European Centre for Disease Prevention and Control (ECDC) reported the age distribution of COVID-19 among children, as of July 2021, in the European Union (EU), European Economic Area (EEA) and the United Kingdom. They found that children made up an increasing proportion of weekly case numbers, with the most noticeable increase among those aged 5-11 years. These findings should be interpreted in light of the proportion of vaccinated adolescents, social mixing patterns by age and adults in those countries at the time [123][135]

Studies from high-income countries have also shown that in some settings, children tend to have more extensive social mixing patterns than adults and consequently more contacts than adults [137]. Thus even though the propensity to transmit may be lower for children, in some settings, they may be contributors to transmission as a consequence of their social mixing patterns, especially if PHSMs have been relaxed [42][126][127][134][141][142][143].

The Omicron variant has resulted in very high levels of incidence in most countries, across all age groups, with higher incidence levels than observed earlier in the pandemic [42]. There is currently limited evidence to suggest a difference in transmission risk of Omicron according to age group, other than that modulated by vaccination, but more data are required. In the context of the Delta and Omicron VOC increased transmission and growth rates have been documented [42].

Figure 1. When should children wear masks?

 World Health Organization

# When should **children** wear masks?

Each country is facing a different situation, as the pandemic evolves, national policies for protective measures should be agile and adjusted as needed.

**Masks are not required**

- Children 5 years of age and under.
- Children with cognitive or respiratory impairments, developmental disorders, disabilities or other specific health conditions that cause them to experience difficulties wearing a mask or who have health conditions that interfere with mask-wearing.
- Children who are doing physical activities, such as running, jumping or playing, since masks may impact their breathing.

**Masks are recommended**

- In areas where there is known or suspected community transmission of SARS-CoV-2, **children ages 6-11 years**:
  - in **indoor** settings where **ventilation is poor or unknown**, even if physical distancing of at least 1 metre can be maintained;
  - in **indoor** settings that have adequate ventilation **when physical distancing of at least 1 metre cannot be maintained**.
- **Adolescents 12 years or older** should follow the same WHO recommendations for mask use as adults.

**Medical masks are recommended**

- **Children with a higher risk\* of severe complications** from COVID-19 should be assessed in consultation with the child's medical provider.
- **Children who have symptoms** of COVID-19 should wear a medical mask at home when they are in shared spaces, as long as they can tolerate it.

\* This includes paediatric patients with underlying noncommunicable diseases (such as diabetes, cardiac disease, chronic lung disease, chronic kidney disease, immunosuppression, HIV, obesity, mental disorders and cancer).

25/03/2022

## Age specific recommendations

## Recommendation for children 5 years of age and under

Conditional recommendation against , Very low certainty evidence



Masks are not required for children 5 years of age and under

Published 7 March 2022

### Practical Info

#### Implementation considerations

As mask use is not recommended in this age group, IPC and public health and social measures should be prioritized to minimize the risk of SARS-CoV-2 transmission.

- Adults and staff working with children should follow national guidelines for vaccination against COVID-19.
- Adequate ventilation\* should be in place and maintained in settings where children are congregating or cared for.
- Adults and staff working with children should wear masks (see WHO recommendations for mask use in adults).
- Adequate sanitation and hygiene requirements and a regimen for environmental cleaning and disinfection should be in place in settings where children congregate or are cared for.
- Children should be taught to perform frequent hand hygiene and respect respiratory etiquette using an age-appropriate approach and materials.

In the event that policymakers decide to adjust the age range for mask recommendations (i.e. children under the age of five years would utilize a mask), relevant settings should have adequate human resources to ensure safe mask use. Adoption of the mask recommendation should include appropriate and consistent supervision by an adult and the ability to ensure mask compliance and adherence, especially if mask-wearing is expected for an extended period. The guiding principles of the best interest of children and a “do no harm” approach should prevail.

*\*For adequate ventilation refer to regional or national institutions or heating, refrigerating and air-conditioning societies implementing ventilation requirements. If recommendations are not in place, a recommended ventilation rate of 10l/s/person should be met (except in healthcare facilities which have specific requirements). For more information, consult [Roadmap to improve and ensure good indoor ventilation in the context of COVID-19](#) [16].*

### Evidence To Decision

#### Benefits and harms

##### Uncertain benefits and harms

The wearing of a well-fitted mask is associated with a decrease in SARS-CoV-2 transmission in the community and provides protective benefits to the individual [29][30][31]. A systematic review on the clinical effectiveness of masks included two RCT and three observational studies in adult populations, which provided some evidence that mask-wearing in the community is associated with decreased risk of COVID-19 infection [30][31][72][79][80][69][70]. The systematic review found inconsistent effects of masks on reducing the risk of influenza-like illness (ILI) in community settings, although a cluster RCT found that hand hygiene and face masks may prevent household transmission of influenza if applied early after symptom onset in an index case [101]. A systematic review evaluating 21 ecological studies in adults reports that mask use is associated with reducing mortality, the incidence of disease, and hospitalization

in the community in the context of COVID-19 [29]. Studies from the United States, Spain, Germany and the United Kingdom looked at the effectiveness of mask use in ages 4-18; and eleven studies reported an association between mask use and decreased COVID-19 incidence in children [144][145][146][147][148][149][150][151][152][153][154]. , These studies were generally observational and ecological with important shortcomings including limited reporting of other infection control measures and exposures.

The systematic review did not find evidence of serious harms with masks in adults in community settings, although bothersome harms were common. Evidence on potential harms, specifically in children aged five years or younger, is limited. Parents who completed an online survey conducted in France reported behavioural and mood changes (e.g. anxiety, sadness, anguish), headaches, speaking difficulties and breathing discomfort attributed to mask-wearing [155]. There is currently no evidence on the long-term impact of mask use on the physical and mental health, development and wellbeing of children.

Given the lack of direct evidence in this age group, evidence was extrapolated from adults. The GDG found that evidence from adults is less applicable (more indirect) to children five and under compared to older children due to lower COVID-19 incidence and severity. Even if masks are associated with the same relative reduction in COVID-19 incidence in children five and under as in adults, the absolute benefits would be smaller due to lower incidence and severity. Furthermore, benefits in children five and under are likely further reduced due to suboptimal adherence.

Additionally, despite the limited/lack of evidence on harms in this age group, there were concerns regarding potential greater harms with regard to childhood development. The GDG, therefore, determined that given the above information, the benefits of mask-wearing in children aged five and under are trivial to none and do not outweigh potential harms.

### Certainty of the Evidence

Very low

The evidence certainty is very low due to the limited evidence in this age group and lower applicability of evidence in adults to this age group compared to older children.

### Values and preferences

Substantial variability is expected or uncertain

The GDG determined that given the close balance of benefits and harms, different preferences (e.g. focusing on potential benefits in terms of reducing infection risk versus focusing on potential developmental harms) could change the decision. Therefore, variability in preferences/values could impact judgments about mask use in this population.

### Resources

No important issues with the recommended alternative

Given that masks are not recommended for this age group, minimal resource implications are anticipated.

### Equity

#### Effect on equity variable

Risk factors that increase the likelihood of contracting COVID-19 include race, ethnicity, and community-level socioeconomic status [156][157].

The GDG assessed effects on equity as uncertain or variable, because masks are not required in this age group, but would depend upon how mask use is implemented. If masks are widely available, using masks could improve equity by reducing the risk of transmission overall, including among socioeconomically disadvantaged groups more impacted by COVID-19. However, there is a need to ensure that lack of access to masks does not negatively impact children (which would decrease equity) and that certain populations (such as disabled individuals) are not adversely impacted.

### Acceptability

There is a significant lack of evidence as to the acceptability of mask use for children in this age group across different contexts<sup>[158][144]</sup>. Additionally, despite limited evidence on harms in this age group, there are concerns regarding potential greater harms with regard to childhood development.

The GDG felt that the acceptability of mask use in children under five years of age is variable.

### Feasibility

The GDG judged that use of masks is less feasible in this age group since it requires more supervision and children may have more difficulty wearing masks for prolonged periods and during certain activities.

### Justification

The GDG determined that benefits of masks in children <5 years did not outweigh harms. This was based on the low certainty evidence and the lower incidence (and severity) of SARS-CoV-2 transmission in this age group relative to older children and adults. The GDG also considered the low acceptability and preference for mask use and agreed that a recommendation for the use of masks for this age group was not appropriate.

Decisions for children under the age of five years to wear masks may be informed by factors such as contact with high-risk individuals, local incidence of COVID-19, ability to adhere to and tolerate mask-wearing, local vaccination rates and parental preferences. There was agreement among the GDG members that in settings where children of this age group are congregating – for example, childcare settings – it is important to adhere to PHSM and IPC measures including adequate ventilation, hand hygiene and environmental hygiene measures, regardless of whether or not masks are used.

## Recommendation for children 6 - 11 years of age

Conditional recommendation for , Low certainty evidence



In areas where there is known or suspected community transmission\* of SARS-CoV- 2, masks are recommended for use in children ages 6-11 years in the following settings:

- in indoor settings where ventilation is known to be poor or cannot be assessed, or the ventilation system is not properly maintained\*\*, regardless of whether physical distancing of at least 1 metre can be maintained,\*\*\*
- in indoor settings that have adequate ventilation\*\* if physical distancing of at least 1 metre cannot be maintained\*\*\*.

\* Details on the levels of community transmission (CT1-CT4) can be found in [Considerations for implementing and adjusting public health and social measures in the context of COVID-19 \[65\]](#). Countries should regularly assess the intensity of spread and health systems capacities at the most localized levels possible.

\*\*For adequate ventilation refer to regional or national institutions or heating, refrigerating and air-conditioning societies implementing ventilation requirements. If regulations are not in place, a recommended ventilation rate of 10l/s/person should be met (except in healthcare facilities which have specific requirements). For more information, consult [Roadmap to improve and ensure good indoor ventilation in the context of COVID-19 \[16\]](#).

\*\*\*Physical distance should be increased beyond 1 metre whenever feasible.

---

Published 7 March 2022

## Practical Info

### Implementation considerations

Countries should regularly assess the intensity of spread and health systems capacities at the most localized levels possible. The assessment should examine the quantitative and qualitative information from available sources and can refer to the situational level (S0-S4) and community transmission (CT) Levels CT1-CT4 as described in [Considerations for implementing and adjusting public health and social measures in the context of COVID-19 \[65\]](#) Additional factors, including population level immunity, will need to be taken into account when setting national and sub-national policies.

Policy and decision-makers are encouraged to ensure the following considerations are addressed when implementing the use of masks in this age group.

- Factors that can influence the decision on implementing the use of masks include the age range in this group, the impact on education and development, routine activities, equity and the general health and wellbeing of children.
- Masks should be made accessible (free of charge) to children in schools, health care settings and any setting where they congregate (e.g. recreational areas), to ensure all children – including those living in households or geographic areas with social vulnerabilities and limited resources – have equitable access. No child should be denied access to these activities for not wearing a mask.
- Efforts should be made to accommodate children who do not have access to masks or are unable to tolerate a mask so they can participate in activities involving face-to-face gatherings. No child should be denied access to these activities for not wearing a mask.
- Routine mask breaks should be implemented when children are expected to wear masks for a longer duration.
- The child's capacity to adhere to correct mask use and availability of appropriate supervision should be addressed, especially in younger children within this age group.

- Age-appropriate communication should aim to help the child understand the purpose and proper use of mask-wearing.
- The design of masks for children should take into consideration the safety and overall quality of the material and ensure a proper fit without compromising breathability, comfort and child-friendliness (appropriate size, colours, patterns).
- Key stakeholders should develop and implement strategies for ensuring that each reusable mask is worn by one child and stored safely, for disposal of soiled masks (e.g. in dedicated bags or containers) and addressing the need for masks to be changed when soiled or wet.
- The use of masks is part of a comprehensive package of preventive measures to reduce transmission including ventilation, physical distance, hand hygiene and respiratory etiquette.

## Evidence To Decision

### Benefits and harms

Substantial net benefits of the recommended alternative

The wearing of a well-fitted mask is associated with a decrease in SARS-CoV-2 transmission in the community and provides protective benefits to the individual [29][30][31]. A systematic review on the clinical effectiveness of masks included two RCT and three observational studies in adult populations that provided some evidence that mask-wearing in the community is associated with decreased risk of COVID-19 infection [30][31]. The systematic review found inconsistent effects of masks on reducing the risk of ILI in community settings, though a cluster RCT found that hand hygiene and face masks may prevent household transmission of influenza if applied early after symptom onset in an index case [101]. A systematic review evaluating 21 ecological studies in adults report that mask use is associated with reducing mortality, the incidence of disease, and hospitalization in the community [29]. Studies from the United States, Spain, Germany and the United Kingdom looked at the effectiveness of mask use in ages 4-18. Ten studies reported an association between mask use and decreased COVID-19 incidence in children. However, these studies were generally observational and ecological with several limitations, including limited reporting of other control measures [145][146][147][148][149][150][151][152][153][154]. Furthermore, two studies of influenza (one RCT and one observational study) found a reduced incidence with mask-wearing in households and school settings [101][104].

The systematic review did not find evidence of serious harms with masks in adults in community settings, although bothersome harms were common. Evidence on potential harms, specifically in children aged 6-11, is limited. Parents who completed an online survey conducted in France - among whom only 9% had children over the age of 11-reported behavioural and mood changes (e.g. anxiety, sadness, anguish), headaches, speaking difficulties and breathing discomfort attributed with mask-wearing [155]. There is currently no evidence on the long-term impact of mask use on the physical and mental health, development and wellbeing of children.

The GDG previously determined that in adults, mask use in community settings is likely associated with a decreased risk of SARS-CoV-2 infections compared with no mask-wearing. The evidence is indirect since it is from adults. Emerging variants such as SARS-CoV-2 B.1.617.2 (Delta) and SARS-CoV-2 B.1.1.529 (Omicron) have been reported to have increased transmissibility [42]. The GDG judged that the benefits in this group are smaller than in adolescents 12 years and older, given lower incidence/severity and reduced adherence (at least in the younger children in this age range).

Evidence on the harms in this age group is also limited. An online survey conducted in France amongst parents of children in a wide age range (<6 years to >11 years) found that parents attributed behavioural change and mood changes (e.g. anxiety, sadness, anguish) headaches, speaking difficulties and breathing discomfort to mask-wearing [155]. However, another study in the United States of America found no apparent adverse biological effects (e.g. impacts on memory, heart rate, oxygen saturation, and emotional state) after mask wearing for at least 30 minutes in elementary school children [159]. There is currently no evidence on the long-term impact of mask use on the physical and mental health, development and wellbeing of children.

The evidence is indirect since it is from adults; the GDG judged that the benefits in this age group are smaller than in adolescents under 12, given lower incidence/severity and reduced adherence (at least in younger children in this age range). Therefore the GDG judged that the benefits of mask-wearing slightly outweigh the harms. Benefits are likely to be larger in situations in which the risk of infection are higher, e.g. poor ventilation and/or unable to physical distance.

### Certainty of the Evidence

Low

There is limited evidence on the benefits and harms of mask-wearing in this age group. Although ecological studies that include children aged 4-18 years have reported an association between mask mandates and a reduced incidence of infection these studies were judged to be low quality, with few studies available from low and middle-income countries [145][146][147][148][149][150][152][153][154][162][163]. Even though this evidence is largely indirect, it was judged by the GDG to have applicability, especially to older children in this group.

### Values and preferences

Substantial variability is expected or uncertain

Substantial variability in preferences, ideas and values is expected regarding the potential outcomes of mask use (prevention of SARS-CoV-2 infection, side effects). Such differences could have an impact on the decision to use masks in this age group.

The GDG determined that given the close balance of benefits and harms, different preferences (e.g., focusing on potential benefits in terms of reducing infection risk versus focusing on potential harms.) could change the decision. Consequently variability in preferences/values could impact judgments about mask use in this population.

### Resources

No important issues with the recommended alternative

There is no formal data available on costs. Given the widespread availability and relatively low costs of non-medical and medical masks, the GDG judged costs and resource availability to be low.

### Equity

#### Effect on equity variable

Risk factors that increase the likelihood of contracting COVID-19 include race, ethnicity, and community-level socioeconomic status [156][157].

The GDG assessed effects on equity as uncertain or variable as it depends on mask use is implemented. If masks are widely available using masks could improve equity by reducing the risk of transmission overall, including among socioeconomically disadvantaged groups more impacted by COVID-19. However, there is a need to ensure that lack of access to masks does not negatively impact children (which would decrease equity) and that certain populations (disabled individuals) are not adversely impacted.

### Acceptability

The limited evidence available indicates variability in the acceptance of masks in children aged 6 to 11. One online study found that parents were generally opposed to children between the ages of 6-10 wearing masks, especially in school settings. Other studies reported that children in this age group demonstrated good adherence to mask-wearing, in particular in school settings [144][152][160].

The GDG decided to make a conditional recommendation despite the low certainty evidence because the benefits of mask-wearing – reduction of SARS-CoV-2 transmission and access to schools – outweigh potential harms, and preferences and values and acceptability generally all favour mask-wearing.

### Feasibility

Adherence is generally feasible in this age group, though there may be some issues in younger children within this

range [144][161].

### Justification

Although there may be a net benefit in mask wearing, this was judged to be small. After reviewing the limited evidence available on the effectiveness of mask use in this age group, a survey was completed by GDG members, among whom 80% voted in favour of a conditional recommendation for mask use. Other factors informing the conditional recommendation were low certainty of evidence, variability in preferences and values that could impact decisions and some variability in acceptability and feasibility.

Settings in which the recommendation applies were also discussed, and members voted 70% in favour of applying the recommendation to indoor settings where ventilation is known to be poor or cannot be assessed or the ventilation system is not adequate and where a distance of at least 1 metre cannot be maintained. The GDG acknowledged the importance of the guiding principles noted earlier, including the right to play and the importance of children continuing to attend school in the context of the COVID-19 pandemic.

### Recommendation for adolescents 12 years of age or older

Strong recommendation for , Low certainty evidence



Adolescents 12 years or older should follow the same WHO [recommendations for mask use as adults](#).

Published 7 March 2022

### Practical Info

#### Implementation consideration

Policy and decision-makers are encouraged to ensure the following considerations are addressed when implementing the use of masks in this age group, irrespective of vaccination status.

- Even where national guidelines apply, additional considerations and adaptations for special settings such as schools, during sports or for children with disabilities or underlying medical conditions will need to be specified.
- Masks should be made accessible free of charge to children in schools, health care settings and any setting where they congregate (such as recreational areas) to ensure all children – including those living in households or geographic areas with social vulnerabilities and limited resources – have equitable access. No child should be denied access to these activities for not wearing a mask.
- Efforts should be made to accommodate children who do not have access to masks or are unable to tolerate a mask so they can participate in activities involving face-to-face gatherings. No child should be denied access to these activities for not wearing a mask.
- Routine mask breaks should be implemented when children are expected to wear masks for a longer duration.
- Age-appropriate communication should aim to help the child understand the purpose and proper use of mask-wearing.
- Key stakeholders should develop and implement strategies for ensuring each reusable mask is worn by one child and stored safely, for disposal of soiled masks (e.g. in dedicated bags or containers) and for addressing the need for masks to be changed when soiled or wet.
- The use of masks is part of a comprehensive package of preventive measures to reduce transmission, including ventilation, physical distance, hand hygiene and respiratory etiquette.

## Evidence To Decision

### Benefits and harms

Small net benefit, or little difference between alternatives

The wearing of a well-fitted mask is associated with a decrease in SARS-CoV-2 transmission in the community and provides protective benefits to the individual [29][30][31]. A systematic review on the clinical effectiveness of masks included two RCT and three observational studies in adult populations that provided some evidence that mask-wearing in the community is associated with decreased risk of COVID-19 infection [30][31]. The systematic review found inconsistent effects of masks on reducing the risk of ILI in community settings, though a cluster RCT found that hand hygiene and face masks may prevent household transmission of influenza if applied early after symptom onset in an index case [101].

A systematic review evaluating 21 ecological studies reports that mask use is associated with reducing mortality, the incidence of disease, and hospitalization in the community [29]. Studies from the United States, Spain, Germany and the United Kingdom looked at the effectiveness of mask use in ages 4-18; twelve studies reported an association between mask use and decreased COVID-19 incidence [145][146][147][148][149][150][151][152][153][154][162][163]. However, these studies were generally observational and ecological with important shortcomings including limited reporting of other infection control measures and exposures.

The systematic review did not find evidence of serious harms with masks in adults in community settings, although bothersome harms were common. Evidence on potential harms specifically in adolescents 12-18 years of age is limited. Parents who completed an online survey conducted in France-among whom only 9% had children over the age of 11-reported behavioural and mood changes (e.g. anxiety, sadness, anguish), headaches, speaking difficulties and breathing discomfort attributed with mask-wearing [155].

The GDG previously determined that in adults, the use of masks in community settings is likely associated with a decreased risk of SARS-CoV-2 infections compared with no mask-wearing. The GDG found that evidence on the use of masks in community settings in adults is likely applicable to adolescents 12 and older due to the similarity in the incidence of SARS-CoV-2 infection (compared with young adults) and ability to adhere to mask-wearing. Emerging variants such as SARS-CoV-2 B.1.617.2 (Delta) and SARS-CoV-2 B.1.1.529 (Omicron) have been reported to have increased transmissibility [42].

The GDG judged the benefits, such as reduced transmission and facilitating increased access to schools/in-person learning, in adolescents to be small but agreed that in the context of the Delta and Omicron variants, the benefits of mask-wearing in the community setting outweigh potential harms.

### Certainty of the Evidence

Low

There is limited evidence on the benefits and harms of mask-wearing in this age group. Although ecological studies that include children aged 4-18 years have reported an association between mask mandates and a reduced incidence of infection these studies were judged to be low quality with few studies available from low and middle-income countries [145][146][147][148][149][150][151][152][153][154][162][163]. Evidence on the effectiveness of masks in adolescents can also be extrapolated from adults. Even though this evidence is indirect, it was judged by the GDG to be more applicable to this age group due to the similarity in incidence and severity of SARS-CoV-2 infection in young adults and adolescents.

### Values and preferences

No substantial variability expected

There is limited data available on adolescents' perception of the value and benefits or harms of wearing masks. Some studies conducted in European settings looking at parental perceptions, showing mixed results but generally favouring mask use in children over the age of 12 [160][164][165]. Given the potential benefits of masks for preventing infections and considering the presence of bothersome but non-serious harms, the GDG determined that differences in values/preference regarding outcomes would not impact the decision to wear masks. This supports a strong recommendation,

despite the low certainty of evidence.

### Resources

No important issues with the recommended alternative

There is no formal data available on costs. Given the widespread availability and relatively low costs non-medical and medical masks, the GDG judged the impact of costs and resource availability to be low.

### Equity

Important issues, or potential issues not investigated

Risk factors that increase the likelihood of contracting COVID-19 include race, ethnicity, and community-level low socioeconomic status [156][157].

The GDG assessed effects on equity as uncertain or variable as it depends on how mask use is implemented. If masks are widely available using masks could improve equity by reducing the risk of transmission overall, including among socioeconomically disadvantaged groups more impacted by COVID-19. However, there is a need to ensure that lack of access to masks does not negatively impact children (which would decrease equity) and that certain populations (such as disabled individuals) are not adversely impacted.

### Acceptability

No important issues with the recommended alternative

This recommendation was assessed by the GDG as likely acceptable in this age group. Studies on the perception of the effectiveness of mask use are limited and generally focused on European countries for children over the age of 10. The GDG considered the limited evidence and discussed knowledge of practice in their respective countries, including the evolution of acceptance of mask use as the pandemic has continued and the emergence of VOC. The GDG agreed that for children over the age of 10 mask-wearing was generally regarded as useful [160][164][165].

### Feasibility

No important issues with the recommended alternative

GDG members noted that masks are widely recommended and used in many contexts throughout the world in this age group. The feasibility of implementing this recommendation was judged to be acceptable and feasible given low concerns about tolerance and likely higher adherence to mask-wearing in older age groups [144].

### Justification

The GDG considered the low certainty of evidence and, although the majority of the evidence was in the adult population, felt it was reasonable to extrapolate from (young) adults. The GDG noted that the benefits of mask use, such as potential reduction in transmission and ability to keep schools functioning, outweighed any potential bothersome harms and considered other factors (not preference-sensitive, low costs, acceptability, feasibility) and believed that this supported a strong recommendation.

## Special populations

#### Good practice statement



Children with cognitive or respiratory impairments, developmental disorders, disabilities\* or other specific health conditions who experience difficulties wearing a mask or have health conditions that interfere with mask-wearing should not be required to wear a mask.

*\* According to the Convention on the Rights of persons with disabilities, children with disabilities "include those who have long-term physical, mental, intellectual or sensory impairments which in interaction with various barriers may hinder their full and effective participation in society on an equal basis" [166].*

Published 7 March 2022

### Practical Info

#### Implementation consideration

- The individual decision for a child to wear a mask should be discussed in consultation with the child's medical provider when possible.
- A safe environment should be created for children who are not able to tolerate a mask, including requirements for caregivers, teachers or other adults interacting with the child to wear a mask when interacting with the child and to be vaccinated against COVID-19 according to national vaccination policies.
- The use of masks with a transparent component may be considered for children with hearing impairment and people who interact with them, where available. These masks should meet approved regulatory standards, if available.

### Justification

The GDG acknowledged that children with several health conditions may experience difficulties or harm while wearing a mask. Despite little direct evidence but considering equity and ethical issues, the GDG determined that a good practice statement was justified.

#### Good practice statement



The use of a medical mask is recommended for children with a higher risk\* of severe complication from COVID-19 but should be assessed in consultation with the child's medical provider.

*\* This includes paediatric patients with underlying non-communicable diseases (for example, diabetes, cardiac disease, chronic lung disease, chronic kidney disease, immunosuppression, obesity, mental disorders and cancer ) and those living with HIV [68].*

Published 7 March 2022

### Justification

The GDG noted that in some low-resource settings there may be challenges for families to access medical masks or have access to a health care provider. It was proposed that in some circumstances it may be more appropriate for caregivers to wear a mask when interacting with the child. In conclusion, the GDG agreed that while there is no direct evidence, a good practice statement was justified due to this population's higher risk of COVID-19 complications.

## Implementation considerations for use of masks in schools

### Implementation consideration

Policy and decision-makers are encouraged to consider the following when implementing mask-wearing by children in school settings.

- Policies should be evidence based, agile and adjusted as needed taking into consideration factors such as changes in transmission intensity, the circulating variant of concern and the capacities for health systems to respond based on the situation.
- No child should be denied access to education because of mask-wearing or the lack of a mask due to low resources or unavailability.
- The views of teachers and educators on risks and time burden required to ensure mask adherence by children should be considered while ensuring that national policies are followed.
- Situations where wearing a mask can significantly interfere with the learning process or have a negative impact on critical school activities such as physical education, or sports and recreation (during which they may reduce ability to breathe comfortably) and meal programmes, require special consideration.
- Specific instructions and supplies should be provided for the availability, safe handling and storage of masks.
- A sufficient supply of appropriate masks should be ensured.
- Masks should not increase social inequalities in access to schools, especially for marginalized communities. No child should be denied access to these activities for not wearing a mask
- Basic water, sanitation, hygiene, ventilation, and space requirements should be met in the school building so that IPC and PHSMs can be implemented.
- If disposable masks are used, a system for waste management of used masks needs to be established to reduce the risk of contaminated masks being disposed of in the classroom and recreational or sports settings.

The recommendations for wearing masks in the different age groups of children in this document supersede those existing in other WHO documents published prior to this update. The following guidance documents can be used to inform policy making and programming for a comprehensive school safety strategy when re-opening or during normal operations in the context of COVID-19:

- [WHO considerations for school-related public health measures in the context of COVID-19](#)
- [WB/WFP/UNESCO/UNICEF framework for school reopening](#)
- [WHO/UNICEF/IFRC Interim Guidance for COVID-19 Prevention and Control in Schools](#)

---

*Published 7 March 2022*

### Justification

GDG members agreed that the recommendations on mask-wearing in this document should be implemented in the context of school settings. They also noted the importance of applying existing public health and social measures and infection prevention and control measures in schools, in addition to mask-wearing.

## Mask use by children authorship, contributors and acknowledgements

*The WHO Health Emergencies and UNICEF Joint Steering group for Masks for Children in the context of COVID-19 (for the update on masks for children, in alphabetical order)*

**Benedetta Allegranzi (WHO/HQ), April Baller (WHO/HQ), Valentine Baltag (WHO/HQ), Anshu Banerjee (WHO/HQ), Anne Detjen (UNICEF), Marjam Esmail (UNICEF), Nathan Ford (WHO/HQ), Nagwa Hasanin (UNICEF), Hannah Hamilton (WHO/HQ),**

**Raoul Kamadjeu** (UNICEF), **Sarah Karmin** (UNICEF), **Jerome Pfaffmann** (UNICEF), **Emma Sacks** (UNICEF), **Wilson Were** (WHO/HQ), **Victoria Willet** (WHO/HQ).

***The joint WHO-UNICEF Guideline Development Group for Masks for children in the context of COVID-19 (November 2021 in alphabetical order)***

**Yewande Alimi** (Africa CDC, Ethiopia), **Jameela Alsalman** (Ministry of Health, Bahrain), **Shelina Bhamani** (Aga Khan University, Pakistan), **Katherine Holland** (Perkins International, United States of America), **Kushlani Jayatilleke** (Sri Jayewardenapura General Hospital, Sri Lanka), **Roberta Petrucci** (Médecins Sans Frontières (MSF), Geneva, Switzerland), **Mathias Pletz** (Jena University Hospital/Friedrich-Schiller-University, The Netherlands), **Fiona Russell** (Department of Paediatrics, The University of Melbourne, Australia), **Marina Salvadori** (Public Health Agency of Canada, Canada), **Paul Anath Tambyah** (National University Hospital, Singapore), **Russell M. Viner** (Faculty of Population Health Sciences, University College London and Royal College of Pediatrics and Child Health, United Kingdom), **Heather Zar** (School of Child and Adolescent Health at the University of Cape Town (UCT), South Africa).

**Methodologist**

**Roger Chou** (Methodologist, Department of Medicine and Department of Medical Informatics and Clinical Epidemiology, Oregon Health & Science University, United States of America)

**Declaration of conflicts of interest**

R. Chou is an author on some of the evidence used to inform some recommendations. However, as a methodologist, he provided guidance to the GDG on methodologic issues and is not a voting member of the GDG. In some meetings, he presented evidence and provided clarification on methods to guide discussions regarding the EtD tables; however, all decisions were made by voting members of the GDG

**Temporary Advisor**

**Dr Fernanda C. Lessa** (US Centers for Disease Control and Prevention, United States of America).

**Declaration of conflicts of interest**

Dr F. Lessa reported she is an employee of the United States CDC who provided funding towards the development of this guideline. After consulting with the WHO Ethics Committee, it was determined Dr Lessa could contribute to discussions as she brings significant technical and field expertise to the discussions but would be recused from voting on recommendations

**External reviewers (for the update on mask use by children, in alphabetical order)**

**Zulfiqar A. Bhutta**, Centre for Global Child Health, The Hospital for Sick Children, Toronto, Canada and Center of Excellence In Women and Child Health, The Aga Khan University Karachi, Pakistan, **Jon Klein**, University of Illinois, United States of America, **Shamez Ladhani**, St. George's University of London, United Kingdom, **Erin Maughan**, College of Health and Human Services, George Mason University, United States of America, **Nina Schwalbe**, Columbia University, United States of America

**Declarations of conflicts of interest**

- *J. Klein* reported receipt of a grant from the International Pediatric Association and UNICEF for contribution of child health and COVID -19 information to pediatric societies. No actions were required.
- *E. Maughan* reported she is also a member of the WHO Technical Advisory Group of Experts on Educational Institutions and COVID-19. No actions were required.
- *Professor F. Russell* declared receipt of funds for research to study school outbreak data and develop mitigation strategies for return to school from the Department of Health, Victoria. No actions were required.

**WHO and UNICEF reviewers (for the update on mask use by children, in alphabetical order)**

**Ida Marie Ameda** (UNICEF), **Astrid Chojnacki** (WHO-WPR), **Landry Cihambanya** (WHO-AFR), **Delphine Sauvageot** (UNICEF), **Aparna Singh Shah** (WHO-SEAR), **Valeska Stempliuk** (WHO-AMR), **Howard Sobel** (WHO-WPR), **Maha Talaat** (WHO-EMR), **Bassim Zayed**, (WHO-EMR)

## Home care for patients

The most up-to-date guidance for "[Home care for patients with suspected or confirmed COVID-19 and management of their contacts](#);

interim guidance” was published 12 August 2020. This guidance is under review and is pending integration into “Infection prevention and control in the context of coronavirus disease (COVID-19): A living guideline”.

## Water, sanitation, hygiene, and waste management

The most up-to-date technical guidance for “Water, sanitation, hygiene, and waste management for SARS-CoV-2, the virus that causes COVID-19” was published 29 July 2020. This guidance is under review and is pending integration into “Infection prevention and control in the context of coronavirus disease (COVID-19): A living guideline”.

## Safe dead body management

The most up-to-date guidance for “Infection prevention and control for the safe management of a dead body in the context of COVID-19: interim guidance” was published 4 September 2020. This guidance is under review and is pending integration into “Infection prevention and control in the context of coronavirus disease (COVID-19): A living guideline”.

## Annexes

### Annex 1. Evidence tables for mask use in the health care setting

This section contains three tables highlighting the application of GRADE to available literature reviewed for mask use in the health care setting.

**Table 1.1** GRADE table for assessment of respirators versus medical mask use in health care settings

<b>Outcome</b>	SARS-CoV-2 infection
<b>Number and type of studies</b>	5 observational studies[54][55][56][57][58]
<b>Consistently</b>	Inconsistent
<b>Precision</b>	No imprecision
<b>Directness</b>	No indirectness
<b>Strength of evidence</b>	Very low
<b>Main findings</b>	<p>Inconsistent findings for N95 vs surgical masks and risk of SARS-Cov-2 infection in health workers.</p> <p>Study 1: OR 1.25 (0.55-2.85) and OR 1.18 (0.86-1.62) [54]</p> <p>Study 2: aOR 7.1(3.6-13.9)[56]</p> <p>Study 3: OR 0.76 (0.63 - 0.92) [55]</p> <p>Study 4: OR 0.60 (0.31 - 1.15) [57]</p> <p>Study 5<sup>^</sup>: PCR+aOR 0.80 (0.64 to 1.00) , seroconversion aOR 0.73 (0.53 - 1.00)[58]</p>

\*All studies were conducted in the pre-delta area except for study 5, which was based on data from June 2020 to February 2021.

<sup>^</sup>Non-peer-reviewed study

**Table 1.2** GRADE table for universal masking versus no universal masking

<b>Outcome</b>	SARS-CoV-2 infection in health workers
<b>Number and type of studies</b>	4 before-after studies[43][44][45][46][47]
<b>Risk of Bias</b>	High
<b>Inconsistently</b>	Not serious
<b>Imprecision</b>	Not serious
<b>Indirectness</b>	Not serious
<b>Quality</b>	Very low
<b>Main findings</b>	Implementation of universal masking temporally associated with reduced incidence of SARS-CoV-2 infection in HCWs

Table 1.3 GRADE table for consistent/always mask use versus inconsistent mask use

<b>Outcome</b>	SARS-CoV-2 infection in health workers
<b>Number and type of studies</b>	6 studies (2 SARS-CoV-2 [170][171], 4 SARS-CoV-1 or MERS-CoV[172][173][174][175])
<b>Risk of Bias</b>	High
<b>Inconsistently</b>	Not serious
<b>Imprecision</b>	Not serious
<b>Indirectness</b>	Serious*
<b>Quality</b>	Very low
<b>Main findings</b>	Consistent/always mask use associated with decreased risk of infection in HCWs vs. inconsistent mask use

\*Most studies were on non-SARS-CoV-2 coronavirus infections; there was insufficient direct evidence from studies of SARS-CoV-2 to determine effects on risk of infection

## Annex 2. Evidence tables for mask use in the community

This section contains a table highlighting the application of GRADE to the available literature reviewed for mask use in community settings.

Table 2.1 GRADE table for assessment of mask versus no mask use in the community setting

<b>Outcome</b>	<b>SARS-CoV-2 infection</b>
<b>Number and type of studies</b>	2 RCTs [69][70] and 10 observational studies [71][72][73][74][75][76][77][78][79][80].
<b>Study limitations</b>	Serious
<b>Consistency</b>	No serious inconsistency
<b>Precision</b>	No serious imprecisions*
<b>Directness</b>	Direct
<b>Strength of evidence</b>	Low-to-moderate
<b>Main findings</b>	2 RCTs: adjusted prevalence ratio 0.90 (95% CI 0.82 to 0.995) for symptomatic SARS-CoV-2 seroprevalence and

	<p>OR 0.82 (95% CI 0.52 to 1.23) for SARS-CoV-2 infection.</p> <p>10 observational studies: OR/HR/RR estimates ranged from 0.04 to 0.86 in 8 studies [71][72][73][74][75][76][77][78][79][80];</p> <p>One additional study of health workers reported an imprecise estimate for mask use outside work (yes vs. no; OR 2.35; 95% CI 0.67 to 8.25) [79] and one study found mask use in a household with an index case of SARS-CoV-2 infection associated with a decreased risk of secondary infection of family members (all family members using mask all the time vs. no family members [OR 0.20; 95% CI 0.07 to 0.60])[80].</p>
--	---

RCT, randomized controlled trial; OR, odds ratio; HR, hazard ratio; RR, relative risk; CI, confidence interval.

\*The RCTs had methodological limitations, including open-label design, attrition, incomplete outcome assessment, variable adherence, and differential recruitment. The RCTs were consistent and were not downgraded for imprecision (due to the very large total sample size [greatly exceeding any optimum information size threshold] with a precise estimate from one of the trials). One trial evaluate a mask recommendation directly; although the other evaluated a mask promotion intervention and did not evaluate mask use or a mask recommendation directly, this resulted in suboptimal uptake of make use and would underestimate the effects of mask use. Therefore, the RCTs were not downgraded for indirectness.

### Annex 3 . Evidence tables for mask use by children.

This section contains two tables highlighting the application of GRADE to available literature reviewed for mask use by children.

**Table 3.1.** GRADE table for assessment of masks versus no mask use in community settings

<b>Outcome</b>	SARS-CoV-2 infection
<b>Number and type of studies</b>	2 RCT and 3 observational studies[31][72][79][80][69][70]
<b>Consistently</b>	Moderate
<b>Precision</b>	Some imprecision*
<b>Directness</b>	Some indirectness*
<b>Strength of evidence</b>	Low-to-moderate
<b>Main findings</b>	<p>RCT1 (cluster): Mask promotion intervention associated with increased mask use and decreased risk of symptomatic SARS-CoV-2 seroprevalence; adjusted prevalence ratio of 0.91, 95% CI 0.82 to 1.00 [69]</p> <p>RCT 2: OR 0.82, 95% CI 0.52 to 1.23 [70]</p> <p>Two observational studies reported inconsistent and imprecise estimates for mask use vs no mask use in community settings outside the home [79][80]. One observational study found mask use by all members of a household or prior to index case illness onset associated with decreased risk of secondary infection vs no mask use [72].</p>

*Note: All studies were conducted in settings without widespread delta variant. Also, ecological studies were not included in this table but consistently found policies requiring masks were associated with decreased risk of SARS-CoV-2 infection.*

\*Of 2 RCTs, one reported an imprecise estimate while the other evaluated an indirect intervention (mask promotion)

**Table 3.2** GRADE assessment of RCTs and observational studies'GRADE assessment of observational and ecological studies on Mask

effectiveness

	Adult Studies	Ecological Studies	Influenza Studies
<b>Outcome</b>	SAR-CoV-2 infection	SARS-CoV-2 infection	SARS-CoV-2 infection
<b>Number of studies</b>	2 RCTs and 3 observational studies [72][79][80][69][70]	13 [145][146][147][148][149][150][151][152][153][154][160][162][163].	1 RCT [101] and 1 observational study [104].
<b>Risk of bias</b>	Moderate	High <sup>2</sup>	Moderate
<b>Consistency</b>	Consistent	Consistent	Consistent
<b>Precision</b>	Some imprecision	Some imprecision	Some imprecision <sup>4</sup>
<b>Directness</b>	Serious indirectness <sup>1</sup>	Serious indirectness <sup>3</sup>	Serious indirectness <sup>5</sup>
<b>Strength of evidence</b>	Low	Very low	Very Low

1 Different population, adult evidence strength rated as moderate. Rated down 1 for children.

2 Studies did not control for the effect of concurrent interventions.

3 Different interventions. Studies did not assess actual mask-wearing or adherence to the intervention

4 RCT outcomes had wide confidence intervals (0.31 - 0.087)

5 Different outcomes were measured. Different population. RCT was a cluster household trial including adults and children. Differences in the intervention: RCT randomized households to facemasks plus 'enhanced hand hygiene' (educational materials provided).

## References

1. Statement on the thirteenth meeting of the International Health Regulations (2005) Emergency Committee regarding the coronavirus disease (COVID-19) pandemic. Geneva: World Health Organization 2022; [Website](#)
2. Chou R, Dana T, Jungbauer R : Update Alert 8: Masks for Prevention of Respiratory Virus Infections, Including SARS-CoV-2, in Health Care and Community Settings. *Annals of internal medicine* 2022;175(9):W108-W109 [Pubmed Journal](#)
3. Coronavirus disease (COVID-19) weekly epidemiological update and weekly operational update [internet]. Geneva: World Health Organization; 2022; [Website](#)
4. Strategic preparedness, readiness and response plan to end the global COVID-19 emergency in 2022. Geneva: World Health Organization 2022; [Website](#)
5. WHO policy brief: Maintaining infection prevention and control measures for COVID-19 in health care facilities, 14 September 2022. Geneva: World Health Organization [Website](#)
6. Version 1.0 of the COVID-19 infection prevention and control living guideline: mask use in community settings. Geneva: World Health Organization; 2021; [Website](#)
7. Mask use in the context of COVID-19: interim guidance, 1 December 2020. Geneva: World Health Organization; 2020; [Website](#)
8. Infection prevention and control in the context of coronavirus disease (COVID-19): a living guideline. Geneva: World Health Organization; 2022; [Website](#)
9. Advice on the use of masks for children in the community in the context of COVID-19. Geneva: World Health Organization; 2020; [Website](#)
10. Infection prevention and control in the context of coronavirus disease (COVID-19): a living guideline, 25 April 2022: updated chapter: mask use, part 1: health care settings. Geneva: World Health Organization 2022;
11. Annex to Infection prevention and control during health care when coronavirus disease (COVID-19) is suspected or confirmed. Geneva: World Health Organization 2021;
12. WHO recommendations on mask use by health workers, in light of the Omicron variant of concern: WHO interim guidelines, 22 December 2021. Geneva: World Health Organization 2021;
13. Schumemann H, Brożek J, Guyatt G, Oxman A : GRADE Handbook. Grading of Recommendations Assessment, Development and Evaluation, Grade Working Group.
14. Guidelines for declaration of interests (WHO experts). Geneva: World Health Organization; 2021; [Website](#)
15. WHO handbook for guideline development, 2nd edition. Geneva: World Health Organization; 2014; [Website](#)
16. Roadmap to improve and ensure good indoor ventilation in the context of COVID-19. Geneva: World Health Organization; 2021; [Website](#)
17. Natural Ventilation for Infection Control in Health-Care Settings. Geneva: World Health Organization; 2009; [Website](#)

18. Infection prevention and control of epidemic-and pandemic prone acute respiratory infections in health care. Geneva: World Health Organization; 2014; [Website](#)
19. Convention on the Rights of the Child: the children's version. New York: UNICEF; 1990; [Website](#)
20. The world health report: 2006: working together for health. Geneva: World Health Organization; 2006; [Website](#)
21. Technical specifications of personal protective equipment for COVID-19. Geneva: World Health Organization; 2020; [Website](#)
22. AFNOR groupe [website]. Saint-Denis: AFNOR International; 2021; [Website](#)
23. ASTM standards & COVID-19 [website]. West Conshohocken (PA): ASTM International; 2021; [Website](#)
24. Directorate General of Drug Administration (DGDA) [website]. Dhaka: Ministry of Health & Family Welfare, Government of the People's Republic of Bangladesh; 2021; [Website](#)
25. European Committee for Standardization [website]. Brussels: Cen; 2021; [Website](#)
26. Siegel JD, Rhinehart E, Jackson M,, Chiarello L, Healthcare Infection Control Practices Advisory Committee. : 2007 Guideline for isolation precautions: preventing transmission of infectious agents in healthcare settings. Atlanta (GA): Centers for Disease Control and Prevention; 2007; [Website](#)
27. Standard precautions for the prevention and control of infections: aide-memoire. Geneva: World Health Organization 2022; [Website](#)
28. Transmission-based precautions for the prevention and control of infections: aide-memoire. Geneva: World Health Organization 2022; [Website](#)
29. Ford N, Holmer HK, Chou R,, Villeneuve PJ, Baller A, et al. : Mask use in community settings in the context of COVID-19: a systematic review of ecological data. *EClinicalMedicine*. 2021;38: 101024 [Pubmed Journal](#)
30. Chou R, Dana T, Jungbauer R,, Weeks C, McDonagh MS : Masks for prevention of respiratory virus infections, including SARS-CoV-2, in health care and community settings: a living rapid review. *Ann Intern Med*. 2020;173(7):542-55. [Pubmed Journal](#)
31. Chou R, Dana T, Jungbauer R : Update alert 6: masks for prevention of respiratory virus infections, including SARS-CoV-2, in health care and community settings. *Ann Intern Med*. 2021;174(9):W68 [Pubmed Journal](#)
32. Chou R, Dana T, Buckley DI,, Selph S, Fu R, Totten AM : Epidemiology of and risk factors for coronavirus infection in health care workers: a living rapid review. *Ann Intern Med*. 2020;173(2):120-36. [Pubmed Journal](#)
33. Chou R, Dana T, Buckley DI,, Selph S, Fu R, Totten AM : Update alert 10: epidemiology of and risk factors for coronavirus infection in health care workers. *Ann Intern Med*. 2022;175(1):W8-W9. [Pubmed Journal](#)
34. Guyatt GH, Alonso-Coello P, Schünemann HJ,, Djulbegovic B, Nothacker M, et al. : Guideline panels should seldom make good practice statements: guidance from the GRADE Working Group. *J Clin Epidemiol*. 2016;80: 3-7. [Pubmed Journal](#)
35. Lotfi T, Hajizadeh A, Moja L,, Akl EA, Piggott T, et al. : A taxonomy and framework for identifying and developing actionable statements in guidelines suggests avoiding informal recommendations. *J Clin Epidemiol*. 2021;141: 161-71. [Pubmed Journal](#)
36. Allegranzi B, Bagheri Nejad S, Combescure C,, Graafmans W, Attar H, et al. : Burden of endemic health-care-associated infection in

developing countries: systematic review and meta-analysis. *Lancet*. 2011;377(9761):228-41. [Pubmed Journal](#)

37. Guidelines on core components of infection prevention and control programmes at the national and acute health care facility level. Geneva: World Health Organization; 2016; [Website](#)
38. Interim practical manual: supporting national implementation of the WHO guidelines on core components of infection prevention and control programmes . Geneva: World Health Organization; 2017; [Website](#)
39. Improving infection prevention and control at the health facility: interim practical manual supporting implementation of the WHO guidelines on core components of infection prevention and control programmes. Geneva: World Health Organization; 2018; [Website](#)
40. Minimum requirements for infection prevention and control programmes. Geneva: World Health Organization; 2019; [Website](#)
41. Core competencies for infection prevention and control professionals. Geneva: World Health Organization; 2020; [Website](#)
42. Enhancing readiness for omicron (B.1.1.529): technical brief and priority actions for member states. Geneva: World Health Organization 2021; [Website](#)
43. Lan F-Y, Christophi CA, Buley J,, Iliaki E, Bruno-Murtha LA, et al. : Effects of universal masking on Massachusetts healthcare workers' COVID-19 incidence. *Occup Med (Lond)*. 2020;70(8):606-9. [Pubmed Journal](#)
44. Seidelman JL, Lewis SS, Advani SD,, Akinboyo IC, Epling C, et al. : Universal masking is an effective strategy to flatten the severe acute respiratory coronavirus virus 2 (SARS-CoV-2) healthcare worker epidemiologic curve. *Infect Control Hosp Epidemiol*. 2020;41(12):1466-7. [Pubmed Journal](#)
45. Wang DD, O'Neill WW, Zervos MJ,, McKinnon JE, Allard D, et al. : Association between implementation of a universal face mask policy for healthcare workers in a health care system and SARS-CoV-2 positivity testing rate in healthcare workers. *J Occup Environ Med*. 2021;63(6):476-81. [Pubmed Journal](#)
46. Wang X, Ferro EG, Zhou G,, Hashimoto D, Bhatt DL : Association between universal masking in a health care system and SARS-CoV-2 positivity among health care workers. *JAMA*. 2020;324(7):703-4. [Pubmed Journal](#)
47. Temkin E, Schwaber MJ, Vaturi A,, Nadir E, Zilber R, et al. : Effect of a national policy of universal masking and uniform criteria for severe acute respiratory coronavirus virus 2 (SARS-CoV-2) exposure on hospital staff infection and quarantine. *Infect Control Hosp Epidemiol*. 2021;3 May 2021: 1-7. [Pubmed Journal](#)
48. Wei J, Li Y : Airborne spread of infectious agents in the indoor environment. *Am J Infect Control*. 2016;44(9 Suppl):S102-8. [Pubmed Journal](#)
49. Liu J, Ma J, Ahmed II,, Varma DK : Effectiveness of a 3D-printed mask fitter in an Ophthalmology setting during COVID-19. *Can J Ophthalmol*. 2021; [Pubmed Journal](#)
50. Leith D, L'Orange C, Volckens J : Quantitative protection factors for common masks and face coverings. *Environ Sci Technol*. 2021;55(5):3136-43. [Pubmed Journal](#)
51. Appendix A to §1910.134—fit testing procedures (mandatory) Part I. OSHA-accepted fit test protocols. Washington (DC): United States Department of Labor; 2004; [Website](#)
52. Tran K, Cimon K, Severn M,, Pessoa-Silva CL, Conly J : Aerosol generating procedures and risk of transmission of acute respiratory infections to healthcare workers: a systematic review. *PLoS One*. 2012;7(4):e35797 [Pubmed Journal](#)

53. van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, et al. : Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *New Engl J Med.* 2020;382(16):1564-7. [Pubmed Journal](#)
54. Fletcher JJ, Feucht EC, Hahn PY, McGoff TN, Dehart DJ, et al. : Healthcare-acquired coronavirus disease 2019 (COVID-19) is less symptomatic than community-acquired disease among healthcare workers. *Infect Control Hosp Epidemiol.* 2021; 1-7. [Pubmed Journal](#)
55. Sims MD, Maine GN, Childers KL, Podolsky RH, Voss DR, et al. : Coronavirus disease 2019 (COVID-19) seropositivity and asymptomatic rates in healthcare workers are associated with job function and masking. *Clin Infect Dis.* 2021;73(Suppl 2):S154-62. [Pubmed Journal](#)
56. Piapan L, De Michieli P, Ronchese F, Rui F, Mauro M, et al. : COVID-19 outbreak in healthcare workers in hospitals in Trieste, north-east Italy. *J Hosp Infect.* 2020;106(3):626-8. [Pubmed Journal](#)
57. Venugopal U, Jilani N, Rabah S, Shariff MA, Jawed M, et al. : SARS-CoV-2 seroprevalence among health care workers in a New York City hospital: a cross-sectional analysis during the COVID-19 pandemic. *Int J Infect Dis.* 2021;102: 63-9. [Pubmed Journal](#)
58. Haller S, Güsewell S, Egger T, Scanferla G, Thoma R : Impact of respirator versus surgical masks on SARS-CoV-2 acquisition in healthcare workers: a prospective multicentre cohort. *Antimicrob Resist Infect Control.* 2022;11(1):27 [Pubmed Journal](#)
59. Jefferson T, Del Mar CB, Dooley L, Ferroni E, Al-Ansary LA, Bawazeer GA : Physical interventions to interrupt or reduce the spread of respiratory viruses. *Cochrane Database Syst Rev.* 2020;11(11):CD006207 [Pubmed Journal](#)
60. Loeb M, Dafoe N, Mahony J, John M, Sarabia A, et al. : Surgical mask vs N95 respirator for preventing influenza among health care workers: a randomized trial. *JAMA.* 2009;302(17):1865-71. [Pubmed Journal](#)
61. Long Y, Hu T, Liu L, Chen R, Guo Q, et al. : Effectiveness of N95 respirators versus surgical masks against influenza: a systematic review and meta-analysis. *J Evid Based Med.* 2020;13(2):93-101. [Pubmed Journal](#)
62. MacIntyre CR, Wang Q, Cauchemez S, Seale H, Dwyer DE, et al. : A cluster randomized clinical trial comparing fit-tested and non-fit-tested N95 respirators to medical masks to prevent respiratory virus infection in health care workers. *Influenza Other Respir Viruses.* 2011;5(3):170-9. [Pubmed Journal](#)
63. MacIntyre CR, Wang Q, Seale H, Yang P, Shi W, et al. : A randomized clinical trial of three options for N95 respirators and medical masks in health workers. *Am J Respir Crit Care Med.* 2013;187(9):960-6. [Pubmed Journal](#)
64. COVID-19 immunization dashboard [internet]. Geneva: World Health Organization; 2021; [Website](#)
65. Considerations for implementing and adjusting public health and social measures in the context of COVID-19. Geneva: World Health Organization; 2021; [Website](#)
66. WHO COVID-19 Case definition. Geneva: World Health Organization 2022; [Website](#)
67. Contact tracing and quarantine in the context of COVID-19: interim guidance, 6 July 2022. Geneva: World Health Organization [Website](#)
68. Living guidance for clinical management of COVID-19: living guidance, 23 November 2021. Geneva: World Health Organization; 2021; [Website](#)
69. Abaluck J, Kwong LH, Styczynski A, Haque A, Kabir MD, et al. : The impact of community masking on COVID-19: a cluster randomized trial in Bangladesh. *Science.* 2022;375(6577): [Pubmed Journal](#)
70. Bundgaard H, Bundgaard JS, Raaschou-Pedersen DET, von Buchwald C, Todsén T, et al. : Effectiveness of adding a mask

recommendation to other public health measures to prevent SARS-CoV-2 infection in Danish mask wearers: a randomized controlled trial. *Ann Intern Med.* 2021;174(3):335-43. [Pubmed Journal](#)

71. Andrejko KL, Pry JM, Myers JF, Fukui N, DeGuzman JL, Openshaw J, et al. : Effectiveness of Face Mask or Respirator Use in Indoor Public Settings for Prevention of SARS-CoV-2 Infection - California, February-December 2021. *MMWR. Morbidity and mortality weekly report* 2022;71(6):212-216 [Pubmed Journal](#)

72. Doung-Ngern P, Suphanchaimat R, Panjangampathana A., Janekrongtham C, Ruampoom D, et al. : Case-control study of use of personal protective measures and risk for SARS-CoV-2 infection, Thailand. *Emerg Infect Dis.* 2020;26(11):2607-16. [Pubmed Journal](#)

73. Gonçalves MR, Dos Reis RCP, Tólio RP, Pellanda LC, Schmidt MI, Katz N, et al. : Social Distancing, Mask Use, and Transmission of Severe Acute Respiratory Syndrome Coronavirus 2, Brazil, April-June 2020. *Emerging infectious diseases* 2021;27(8):2135-2143 [Pubmed Journal](#)

74. Lio CF, Cheong HH, Lei CI, Lo IL, Yao L, Lam C, et al. : Effectiveness of personal protective health behaviour against COVID-19. *BMC public health* 2021;21(1):827 [Pubmed Journal](#)

75. Rebmann T, Loux TM, Arnold LD, Charney R, Horton D, Gomel A : SARS-CoV-2 Transmission to Masked and Unmasked Close Contacts of University Students with COVID-19 - St. Louis, Missouri, January-May 2021. *MMWR. Morbidity and mortality weekly report* 2021;70(36):1245-1248 [Pubmed Journal](#)

76. Sharif N, Alzahrani KJ, Ahmed SN, Opu RR, Ahmed N, Talukder A, et al. : Protective measures are associated with the reduction of transmission of COVID-19 in Bangladesh: A nationwide cross-sectional study. *PloS one* 2021;16(11):e0260287 [Pubmed Journal](#)

77. Sugimura M, Chimed-Ochir O, Yumiya Y, Ohge H, Shime N, Sakaguchi T, et al. : The Association between Wearing a Mask and COVID-19. *International journal of environmental research and public health* 2021;18(17): [Pubmed Journal](#)

78. Tjaden AH, Gibbs M, Runyon M, Weintraub WS, Taylor YJ, Edelstein SL, et al. : Association between Self-reported Masking Behavior and SARS-CoV-2 Infection Wanes from Pre-Delta to Omicron-Predominant Periods - North Carolina COVID-19 Community Research Partnership (NC-CCRP). *American journal of infection control* 2022; [Pubmed Journal](#)

79. van den Broek-Altenburg EM, Atherly AJ, Diehl SA., Gleason KM, Hart VC, et al. : Jobs, housing, and mask wearing: cross-sectional study of risk factors for COVID-19. *JMIR Public Health Surveill.* 2021;7(1): [Pubmed Journal](#)

80. Wang YU, Tian H, Zhang L., Zhang M, Guo D, et al. : Reduction of secondary transmission of SARS-CoV-2 in households by face mask use, disinfection and social distancing: a cohort study in Beijing, China. *BMJ Glob Health.* 2020;5(5): [Pubmed Journal](#)

81. Levine Z, Earn DJD : Face masking and COVID-19: potential effects of variolation on transmission dynamics. *Journal of the Royal Society, Interface* 2022;19(190):20210781 [Pubmed Journal](#)

82. Hale T, Angrist N, Goldszmidt R, Kira B, Petherick A, Phillips T, et al. : A global panel database of pandemic policies (Oxford COVID-19 Government Response Tracker). *Nature human behaviour* 2021;5(4):529-538 [Pubmed Journal](#)

83. Face masks and coverings for the general public: Behavioural knowledge, effectiveness of cloth coverings and public messaging. *The Royal Society* 2020; [Website](#)

84. Tsang PM, Prost A : Boundaries of solidarity: a meta-ethnography of mask use during past epidemics to inform SARS-CoV-2 suppression. *BMJ global health* 2021;6(1): [Pubmed Journal](#)

85. Roberge RJ : Face shields for infection control: a review. *J Occup Environ Hyg.* 2016;13(4):235-42. [Pubmed Journal](#)

86. Machida M, Nakamura I, Saito R., Nakaya T, Hanibuchi T, et al. : Incorrect use of face masks during the current COVID-19 pandemic

among the general public in Japan. *Int J Environ Res Public Health*. 2020;17(18):6484 [Pubmed Journal](#)

87. Milton DK, Fabian MP, Cowling B,, Grantham ML, McDevitt JJ : Influenza virus aerosols in human exhaled breath: particle size, culturability, and effect of surgical masks. *PLoS Pathog*. 2013;9(3): [Pubmed Journal](#)

88. Chen Y-J, Qin G, Chen J,, Xu J-L, Feng D-Y, et al. : Comparison of face-touching behaviors before and during the coronavirus disease 2019 pandemic. *JAMA Netw Open*. 2020;3(7):e2016924 [Pubmed Journal](#)

89. Shiraly R, Shayan Z, McLaws M-L : Face touching in the time of COVID-19 in Shiraz, Iran. *Am J Infect Control*. 2020;48(12):1559-61. [Pubmed Journal](#)

90. Betsch C, Korn L, Sprengholz P,, Felgendreff L, Eitze S, Schmid P : Social and behavioral consequences of mask policies during the COVID-19 pandemic. *Proc Natl Acad Sci U S A* . 2020;117(36):21851-3. [Pubmed Journal](#)

91. Cowling BJ, Ali ST, Ng T,, Tsang TK, Li JCM, Fong MW : Impact assessment of non-pharmaceutical interventions against coronavirus disease 2019 and influenza in Hong Kong: an observational study. *Lancet Public Health*. 2020;5(5):e279-88. [Pubmed Journal](#)

92. Bakhit M, Krzyzaniak N, Scott A,, Clark J, Glasziou P, Del Mar C : Downsides of face masks and possible mitigation strategies: a systematic review and meta-analysis. *BMJ Open*. 2021;11(2):e044364 [Pubmed Journal](#)

93. Kunstler B, Newton S, Hill H, Ferguson J, Hore P, Mitchell BG, et al. : P2/N95 respirators & surgical masks to prevent SARS-CoV-2 infection: Effectiveness & adverse effects. *Infection, disease & health* 2022;27(2):81-95 [Pubmed Journal](#)

94. Kisielinski K, Giboni P, Prescher A, Klosterhalfen B, Graessel D, Funken S, et al. : Is a Mask That Covers the Mouth and Nose Free from Undesirable Side Effects in Everyday Use and Free of Potential Hazards?. *International journal of environmental research and public health* 2021;18(8): [Pubmed Journal](#)

95. Balestracci B, La Regina M, Di Sessa D, Mucci N, Angelone FD, D'Ecclesia A, et al. : Patient safety implications of wearing a face mask for prevention in the era of COVID-19 pandemic: a systematic review and consensus recommendations. *Internal and emergency medicine* 2022; [Pubmed Journal](#)

96. Thatiparthi A, Liu J, Martin A, Wu JJ : Adverse Effects of COVID-19 and Face Masks: A Systematic Review. *The Journal of clinical and aesthetic dermatology* 2021;14(9 Suppl 1):S39-S45 [Pubmed](#)

97. Giacalone S, Minuti A, Spigariolo C,, Passoni E, Nazzaro G : Facial dermatoses in the general population due to wearing of personal protective masks during the COVID-19 pandemic: first observations after lockdown. *Clin Exp Dermatol*. 2021;46(2):368-9. [Pubmed Journal](#)

98. Matusiak Ł, Szepietowska M, Krajewski P,, Białynicki-Birula R, Szepietowski JC : Inconveniences due to the use of face masks during the COVID-19 pandemic: a survey study of 876 young people. *Dermatol Ther*. 2020;33(4):e13567 [Pubmed Journal](#)

99. Hüfner K, Hofer A, Sperner-Unterweger B : On the difficulties of building therapeutic relationships when wearing face masks. *J Psychosom Res*. 2020;138(110226):110226 [Pubmed Journal](#)

100. Crume B : The silence behind the mask: my journey as a deaf pediatric resident amid a pandemic. *Acad Pediatr*. 2020;21(1):1-2. [Pubmed Journal](#)

101. Cowling BJ, Chan K-H, Fang V,, Cheng CKY, Fung ROP, et al. : Facemasks and hand hygiene to prevent influenza transmission in households: a cluster randomized trial. *Ann Intern Med*. 2009;151(7):437-46. [Pubmed Journal](#)

102. Allison MA, Guest-Warnick G, Nelson D,, Pavia AT, Srivastava R, et al. : Feasibility of elementary school children's use of hand gel and facemasks during influenza season. *Influenza Other Respir Viruses*. 2010;4(4):223-9. [Pubmed Journal](#)

103. Canini L, Andréoletti L, Ferrari P, D'Angelo R, Blanchon T, et al. : Surgical mask to prevent influenza transmission in households: a cluster randomized trial. *PLoS One*. 2010;5(11): [PubMed Journal](#)
104. Uchida M, Kaneko M, Hidaka Y, Yamamoto H, Honda T, et al. : Effectiveness of vaccination and wearing masks on seasonal influenza in Matsumoto City, Japan, in the 2014/2015 season: An observational study among all elementary schoolchildren. *Prev Med Rep*. 2017;5: 86-91. [PubMed Journal](#)
105. Zand AD, Heir AV : Environmental impacts of new coronavirus outbreak in Iran with an emphasis on waste management sector. *J Mater Cycles Waste Manag*. 2021;23(1):240-7. [PubMed Journal](#)
106. Global analysis of health care waste in the context of COVID-19. Geneva: World Health Organization [Website](#)
107. Coronavirus disease (COVID-19) advice for the public: when and how to use masks [internet]. Geneva: World Health Organization; 2020; [Website](#)
108. Scott N, Saul A, Spelman T, Stoope M, Pedrana A, et al. : The introduction of a mandatory mask policy was associated with significantly reduced COVID-19 cases in a major metropolitan city. *PLoS One*. 2021;16(7): [PubMed Journal](#)
109. Aydin O, Emon B, Cheng S, Hong L, Chamorro LP, Saif MTA : Performance of fabrics for home-made masks against the spread of COVID-19 through droplets: a quantitative mechanistic study. *Extreme Mech Lett*. 2020;40. [PubMed Journal](#)
110. Fischer EP, Fischer MC, Grass D, Henrion I, Warren WS, Westman E : Low-cost measurement of face mask efficacy for filtering expelled droplets during speech. *Sci Adv*. 2020;6(36): [PubMed Journal](#)
111. Bull FC, Al-Ansari SS, Biddle S, Borodulin K, Buman MP, Cardon G : World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med*. 2020;54(24):1451-62. [PubMed Journal](#)
112. Wong AY-Y, Ling SK-K, Louie LH-T, Law GY-K, So RC-H, Lee DC-W, et al. : Impact of the COVID-19 pandemic on sports and exercise. *Asia-Pacific journal of sports medicine, arthroscopy, rehabilitation and technology* 2020;22 39-44 [PubMed Journal](#)
113. Fikenzer S, Uhe T, Lavall D, Rudolph U, Falz R, Busse M : Effects of surgical and FFP2/N95 face masks on cardiopulmonary exercise capacity. *Clin Res Cardiol*. 2020;109(12):1522-30. [PubMed Journal](#)
114. Kyung SY, Kim Y, Hwang H, Park J-W, Jeong SH : Risks of N95 face mask use in subjects with COPD. *Respir Care*. 2020;65(5):658-64. [PubMed Journal](#)
115. Yang G-Z, Kelley E, Darzi A : Patients' safety for global health. *Lancet*. 2010;377(9769):886-7. [PubMed Journal](#)
116. Lee HP, Wang DY : Objective assessment of increase in breathing resistance of N95 respirators on human subjects. *Ann Occup Hyg*. 2011;55(8):917-21. [PubMed Journal](#)
117. Harber P, Santiago S, Bansal S, Liu Y, Yun D, Wu S : Respirator physiologic impact in persons with mild respiratory disease. *J Occup Environ Med*. 2010;52(2):155-62. [PubMed Journal](#)
118. Person E, Lemercier C, Royer A, Reychler G : Effet du port d'un masque de soins lors d'un test de marche de six minutes chez des sujets sains [Effect of a surgical mask on six minute walking distance]. *Rev Mal Respir*. 2018;35(3):264-8. [PubMed Journal](#)
119. Li Y, Tokura H, Guo Y, Wong ASW, Wong T, et al. : Effects of wearing N95 and surgical facemasks on heart rate, thermal stress and subjective sensations. *Int Arch Occup Environ Health*. 2005;78(6):501-9. [PubMed Journal](#)

120. Hopkins SR, Dominelli PB, Davis C., Guenette JA, Luks AM, et al. : Face masks and the cardiorespiratory response to physical activity in health and disease. *Ann Am Thorac Soc.* 2021;18(3):399-407. [Pubmed Journal](#)
121. Lee LY-K, Lam EP-W, Chan C-K, Chan S-Y, Chiu M-K, Chong W-H, et al. : Practice and technique of using face mask amongst adults in the community: a cross-sectional descriptive study. *BMC public health* 2020;20(1):948 [Pubmed Journal](#)
122. Bialek S, Gierke R, Hughes M., McNamara LA, Pilishvili T, Skoff T : Coronavirus disease 2019 in children - United States, February 12-April 2, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69(14):422-6. [Pubmed Journal](#)
123. COVID-19 in children and the role of school settings in transmission – second update. Stockholm: European Centre for Disease Prevention and Control; 2021; [Website](#)
124. Fontanet A, Tondeur L, Grant R., Temmam S, Madec Y, et al. : SARS-CoV-2 infection in schools in a northern French city: a retrospective serological cohort study in an area of high transmission, France, January to April 2020. *Euro Surveill.* 2021;26(15):2001695 [Pubmed Journal](#)
125. Goldstein E, Lipsitch M, Cevik M : On the effect of age on the transmission of SARS-CoV-2 in households, schools, and the community. *J Infect Dis.* 2020;223(3):362-9. [Pubmed Journal](#)
126. Ladhani SN, Amin-Chowdhury Z., Davies H., Aiano F, Hayden I, et al. : COVID-19 in children: analysis of the first pandemic peak in England. *Arch Dis Child.* 2020;105(12):1180-5. [Pubmed Journal](#)
127. Ladhani SN, Ireland G, Baawuah F., Beckmann J, Okike IO, et al. : SARS-CoV-2 infection, antibody positivity and seroconversion rates in staff and students following full reopening of secondary schools in England: a prospective cohort study, September–December 2020. *EClinicalMedicine.* 2021;37: 100948 [Pubmed Journal](#)
128. Li X, Xu W, Dozier M., He Y, Kirolos A, et al. : The role of children in the transmission of SARS-CoV2: updated rapid review. *J Glob Health.* 2020;10(2):021101 [Pubmed Journal](#)
129. Ludvigsson JF : Children are unlikely to be the main drivers of the COVID-19 pandemic: a systematic review. *Acta Paediatr.* 2020;109(8):1525-30. [Pubmed Journal](#)
130. Lu Y, Li Y, Deng W., Liu M, He Y, et al. : Symptomatic infection is associated with prolonged duration of viral shedding in mild coronavirus disease 2019: a retrospective study of 110 children in Wuhan. *Pediatr Infect Dis J.* 2020;39(7):e95-9. [Pubmed Journal](#)
131. National COVID-19 surveillance reports, including weekly summary of findings monitored through various COVID-19 surveillance systems [internet]. London: Public Health England; 2020; [Website](#)
132. Stringhini S, Wisniak A, Piumatti G., Azman AS, Lauer SA, et al. : Seroprevalence of anti-SARS-CoV-2 IgG antibodies in Geneva, Switzerland (SEROCoV-POP): a population-based study. *Lancet.* 2020;396(10247):313-9. [Pubmed Journal](#)
133. Torres JP, Piñera C, De La Maza V., Lagomarcino AJ, Simian D, et al. : Severe acute respiratory syndrome coronavirus 2 antibody prevalence in blood in a large school community subject to a coronavirus disease 2019 outbreak: a cross-sectional study. *Clin Infect Dis.* 2020;73(2):e458-65. [Pubmed Journal](#)
134. Viner RM, Mytton OT, Bonell C., Melendez-Torres GJ, Ward J, et al. : Susceptibility to SARS-CoV-2 infection among children and adolescents compared with adults: a systematic review and meta-analysis. *JAMA Pediatr.* 2021;175(2):143-56. [Pubmed Journal](#)
135. Chadeau-Hyam M, Wang H, Eales O., Haw D, Bodinier B, et al. : REACT-1 study round 14: High and increasing prevalence of SARS-CoV-2 infection among school-aged children during September 2021 and vaccine effectiveness against infection in England. *medRxiv [preprint].* 2021; [Journal](#)

136. Elliot P, Bodineier B, Ealse O,, Wang H, Haw D : Rapid increase in Omicron infections in England during December 2021: REACT-1 study. London: Imperial College London; 2021; [Website](#)
137. Munday J, Jarvis C, Gimma A,, Wong K, Van Zandvoort K, et al. : Contact matrices estimated from the CoMix social contact survey. London: London School of Hygiene and tropical Medicine; 2021; [Website](#)
138. Kleynhans J, Tempia S, Wolter N,, von Gottberg A, Bhiman J, et al. : SARS-CoV-2 seroprevalence in a rural and urban household cohort during first and second waves of infections, South Africa, July 2020–March 2021. *Emerg Infect Dis.* 2021;27(12):3020-9. [Pubmed Journal](#)
139. Weekly epidemiological update on COVID-19; 19-October-2021 [internet]. Geneva: World Health Organization; 2021; [Website](#)
140. Real-time assessment of community transmission findings [internet]. London: Imperial College London; 2021; [Website](#)
141. COVID-19 SeroHub [internet]. Washington (DC): National Institutes of Health; 2021; [Website](#)
142. Thompson HA, Mousa A, Dighe A,, Fu H, Arnedo-Pena A, et al. : Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) setting-specific transmission rates: a systematic review and meta-analysis. *Clin Infect Dis.* 2021;73(3):e754-64. [Pubmed Journal](#)
143. Zhu Y, Bloxham CJ, Hulme KD,, Sinclair JE, Tong ZWM, et al. : A meta-analysis on the role of children in severe acute respiratory syndrome coronavirus 2 in household transmission clusters. *Clin Infect Dis.* 2021;72(12):e1146-53. [Pubmed Journal](#)
144. Mickells GE, Figueroa J, West K,, Wood A, McElhanon BO : Adherence to masking requirement during the COVID-19 pandemic by early elementary school children. *J Sch Health.* 2021;91(7):555-61. [Pubmed Journal](#)
145. Budzyn SE, Panaggio MJ, Parks S,, Papazian M, Magid J, Barrios LC : Pediatric COVID-19 cases in counties with and without school mask requirements—United States, July 1-September 4, 2021. *MMWR Morb Mortal Wkly Rep.* 2021;70 1377-8. [Pubmed Journal](#)
146. Chernozhukov V, Kasahara H, Schrimpf P : The association of opening K–12 schools with the spread of COVID-19 in the United States: county-level panel data analysis. *Proc Natl Acad Sci U S A.* 2021;118(42):e2103420118 [Pubmed Journal](#)
147. Doyle T, Kendrick K, Troelstrup T,, Gumke M, Edwards J, et al. : COVID-19 in primary and secondary school settings during the first semester of school reopening—Florida, August-December 2020. *MMWR Morb Mortal Wkly Rep.* 2021;70(12):437-41. [Pubmed Journal](#)
148. Falk A, Benda A, Falk P,, Steffen S, Wallace Z, Hoeg TB : COVID-19 cases and transmission in 17 K-12 schools—Wood County, Wisconsin, August 31-November 29, 2020. *MMWR Morb Mortal Wkly Rep.* 2021;70(4):136-40. [Pubmed Journal](#)
149. Gettings J, Czarnik M, Morris E,, Haller E, Thompson AM, et al. : Mask use and ventilation improvements to reduce COVID-19 incidence in elementary schools—Georgia, November 16–December 11, 2020. *MMWR Morb Mortal Wkly Rep.* 2021;70 779-84. [Pubmed Journal](#)
150. Jordan I, de Sevilla MF, Fumado V,, Bassat Q, Bonet-Carne E, et al. : Transmission of SARS-CoV-2 infection among children in summer schools applying stringent control measures in Barcelona, Spain. *Clin Infect Dis.* 2021;74(1):66-73. [Pubmed Journal](#)
151. McCullough JM, Dale AP, Gue M,, Eller B, Cullen T, Scott SE : Association between K–12 school mask policies and school-associated COVID-19 outbreaks— Maricopa and Pima Counties, Arizona, July–August 2021. *MMWR Morb Mortal Wkly Rep.* 2021;70(39):1372-3. [Pubmed Journal](#)
152. Nelson SB, Dugdale CM, Bilinski A,, Cosar D, Pollock NR, Ciaranello A : Prevalence and risk factors for in-school transmission of SARS-CoV-2 in Massachusetts K-12 public schools, 2020-2021. *medRxiv [preprint].* 2021; [Journal](#)

153. Yuan H, Reynolds C, Ng S., Yang W : Factors affecting the transmission of SARS-CoV-2 in school settings. *Influenza Other Respir Viruses*. 2022; [Pubmed Journal](#)
154. Zimmerman KO, Akinboyo IC, Brookhart M., Boutzoukas AE, McGann KA, et al. : Incidence and secondary transmission of SARS-CoV-2 infections in schools. *Pediatrics*. 2021;147(4):e2020048090 [Pubmed Journal](#)
155. Assathiany R, Salinier C, Béchet S., Dolard C, Kochert F, et al. : Face masks in young children during the COVID-19 pandemic: parents' and pediatricians' point of view. *Front Pediatr*. 2021 2021;9(579):676718 [Journal](#)
156. Lo C-H, Nguyen LH, Drew DA., Warner ET, Joshi AD, et al. : Race, ethnicity, community-level socioeconomic factors, and risk of COVID-19 in the United States and the United Kingdom. *EClinicalMedicine*. 2021;38 101029 [Pubmed Journal](#)
157. Sze S, Pan D, Nevill CR., Gray LJ, Martin CA, et al. : Ethnicity and clinical outcomes in COVID-19: a systematic review and meta-analysis. *EClinicalMedicine*. 2020;29 100630 [Pubmed Journal](#)
158. Aronu AE, Chinawa JM, Nduagubam OC., Ossai EN, Chinawa AT, Igwe WC : Maternal perception of masking in children as a preventive strategy for COVID-19 in Nigeria: A multicentre study. *PLoS One*. 2020;15(11): [Pubmed Journal](#)
159. Smith J, Culler A, Scanlon K : Impacts of blood gas concentration, heart rate, emotional state, and memory in school-age children with and without the use of facial coverings in school during the COVID-19 pandemic. *FASEB J*. 2021;35(S1): [Journal](#)
160. Betsch C., Korn L., Felgendreff L., Eitze S., Thaiss H. : School opening during the SARS-CoV-2 pandemic: Public acceptance of wearing fabric masks in class. *Public Health Pract (Oxf)* 2021;2 100115 [Pubmed Journal](#)
161. Chen X, Ran LI, Liu Q., Hu Q, Du X, Tan X : Hand hygiene, mask-wearing behaviors and its associated factors during the COVID-19 epidemic: a cross-sectional study among primary school students in Wuhan, China. *Int J Environ Res Public Health*. 2020;17(8):2893 [Pubmed Journal](#)
162. Theuring S, Thielecke M, van Loon W., Hommes F, Hülso C, et al. : SARS-CoV-2 infection and transmission in school settings during the second COVID-19 wave: a cross-sectional study, Berlin, Germany, November 2020. *Euro Surveill*. 2021;26(34):2100184 [Pubmed Journal](#)
163. Volpp KG, Kraut BH, Ghosh S., Neatherlin J : Minimal SARS-CoV-2 transmission after implementation of a comprehensive mitigation strategy at a school—New Jersey, August 20–November 27, 2020. *MMWR Morb Mortal Wkly Rep*. 2021;70(11):377–81. [Pubmed Journal](#)
164. Ammann P, Ulyte A, Haile SR., Puhan MA, Kriemler S, Radtke T : Perceptions towards mask use in school children during the SARS-CoV-2 pandemic: the Ciao Corona study. *medRxiv [preprint]*. 2021; [Journal](#)
165. Ayran G, Köse S, Sarılioğlu A., Çelebioğlu A : Hand hygiene and mask-wearing behaviors and the related factors during the COVID 19 pandemic: a cross-sectional study with secondary school students in Turkey. *J Pediatr Nurs*. 2022;62: 98-105. [Pubmed Journal](#)
166. Convention on the Rights of Persons with Disabilities. Article 2. New York (NY): United Nations; 2008; [Website](#)
167. Infection prevention and control (IPC) in health-care facilities in the event of a surge or resurgence in cases of COVID-19. Geneva: World Health Organization
168. Lustig SR, Biswakarma JJH, Rana D, Tilford SH, Hu W, Su M, et al. : Effectiveness of Common Fabrics to Block Aqueous Aerosols of Virus-like Nanoparticles. *ACS nano* 2020;14(6):7651-7658 [Pubmed Journal](#)
169. Lee S-A, Hwang D-C, Li H-Y, Tsai C-F, Chen C-W, Chen J-K : Particle Size-Selective Assessment of Protection of European Standard FFP Respirators and Surgical Masks against Particles-Tested with Human Subjects. *Journal of healthcare engineering* 2016;2016 [Pubmed Journal](#)

170. Akinbami LJ, Vuong N, Petersen LR, Sami S, Patel A, Lukacs SL, et al. : SARS-CoV-2 Seroprevalence among Healthcare, First Response, and Public Safety Personnel, Detroit Metropolitan Area, Michigan, USA, May-June 2020. *Emerging infectious diseases* 2020;26(12):2863-2871 [Pubmed Journal](#)
171. Davido B, Gautier S, Riom I, Landowski S, Lawrence C, Thiebaut A, et al. : The first wave of COVID-19 in hospital staff members of a tertiary care hospital in the greater Paris area: A surveillance and risk factors study. *International journal of infectious diseases : IJID : official publication of the International Society for Infectious Diseases* 2021;105 172-179 [Pubmed Journal](#)
172. Loeb M, McGeer A, Henry B, Ofner M, Rose D, Hlywka T, et al. : SARS among critical care nurses, Toronto. *Emerging infectious diseases* 2004;10(2):251-5 [Pubmed](#)
173. Nishiyama A, Wakasugi N, Kirikae T, Quy T, Ha LD, Ban VV, et al. : Risk factors for SARS infection within hospitals in Hanoi, Vietnam. *Japanese journal of infectious diseases* 2008;61(5):388-90 [Pubmed](#)
174. Alraddadi BM, Al-Salmi HS, Jacobs-Slifka K, Slayton RB, Estivariz CF, Geller AI, et al. : Risk Factors for Middle East Respiratory Syndrome Coronavirus Infection among Healthcare Personnel. *Emerging infectious diseases* 2016;22(11):1915-1920 [Pubmed Journal](#)
175. Lau JTF, Fung KS, Wong TW, Kim JH, Wong E, Chung S, et al. : SARS transmission among hospital workers in Hong Kong. *Emerging infectious diseases* 2004;10(2):280-6 [Pubmed](#)